

Datasheet for ABIN7551534

POLR2L Protein (AA 1-67) (Fc Tag)



Overview

Quantity:	1 mg			
Target:	POLR2L			
Protein Characteristics:	AA 1-67			
Origin:	Human			
Source:	HEK-293 Cells			
Protein Type:	Recombinant			
Purification tag / Conjugate:	This POLR2L protein is labelled with Fc Tag.			
Application:	Western Blotting (WB), SDS-PAGE (SDS)			
Product Details				
Purpose:	Custom-made recombinat POLR2L Protein expressed in mammalien cells.			
Sequence:	MIIPVRCFTC GKIVGNKWEA YLGLLQAEYT EGDALDALGL KRYCCRRMLL AHVDLIEKLL			
	NYAPLEK Sequence without tag. The proposed Purification-Tag is based on experiences			
	with the expression system, a different complexity of the protein could make another tag			
	necessary. In case you have a special request, please contact us.			
Characteristics:	Key Benefits:			
	 Made to order protein - from design to production - by highly experienced protein experts. Protein expressed in mammalien cells and purified in one-step affinity chromatography The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins. State-of-the-art algorithm used for plasmid design (Gene synthesis). 			

This protein is a made-to-order protein and will be made for the first time for your order. Our

Product Details		
	experts in the lab try to ensure that you receive soluble protein.	
	If you are not interested in a full length protein, please contact us for individual protein fragments.	
	The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.	
Purity:	> 90 % as determined by Bis-Tris Page, Western Blot	
Grade:	custom-made	
Target Details		
Target:	POLR2L	
Alternative Name:	POLR2L (POLR2L Products)	
Background:	DNA-directed RNA polymerases I, II, and III subunit RPABC5 (RNA polymerases I, II, and III subunit ABC5) (DNA-directed RNA polymerase III subunit L) (RNA polymerase II 7.6 kDa subunit) (RPB7.6) (RPB10 homolog),FUNCTION: DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. Common component of RNA polymerases I, II and III which synthesize ribosomal RNA precursors, mRNA precursors and many functional non-coding RNAs, and a small RNAs, such as 5S rRNA and tRNAs, respectively. {ECO:0000250, ECO:0000250 UniProtKB:P22139,	

Molecular Weight:	7.6 kDa
	ECO:0000269 PubMed:36271492, ECO:0000269 PubMed:9852112}.
	ECO:0000269 PubMed:34671025, ECO:0000269 PubMed:34887565,
	ECO:0000269 PubMed:27193682, ECO:0000269 PubMed:30190596,
	ECO:0000269 PubMed:16809778, ECO:0000269 PubMed:20413673,
	as 5S rRNA and tRNAs, respectively. {ECO:0000250, ECO:0000250 UniProtKB:P22139,
	precursors, mRNA precursors and many functional non-coding RNAs, and a small RNAs, such
	Common component of RNA polymerases I, II and III which synthesize ribosomal RNA
	transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates.
	subunit) (RPB7.6) (RPB10 homolog),FUNCTION: DNA-dependent RNA polymerase catalyzes the

Molecular Weight:	7.6 kDa	
UniProt:	P62875	

Regulatory RNA Pathways Pathways:

Application Details

In addition to the applications listed above we expect the protein to work for functional studies Application Notes: as well. As the protein has not been tested for functional studies yet we cannot offer a

Application Details

	guarantee though.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months