

Datasheet for ABIN7553208

Cyclin C Protein (CCNC) (AA 1-283) (His tag)



Overview

Quantity:	1 mg
Target:	Cyclin C (CCNC)
Protein Characteristics:	AA 1-283
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Cyclin C protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

Product Details	
Purpose:	Custom-made recombinat CCNC Protein expressed in mammalien cells.
Sequence:	MAGNFWQSSH YLQWILDKQD LLKERQKDLK FLSEEEYWKL QIFFTNVIQA LGEHLKLRQQ
	VIATATVYFK RFYARYSLKS IDPVLMAPTC VFLASKVEEF GVVSNTRLIA AATSVLKTRF
	SYAFPKEFPY RMNHILECEF YLLELMDCCL IVYHPYRPLL QYVQDMGQED MLLPLAWRIV
	NDTYRTDLCL LYPPFMIALA CLHVACVVQQ KDARQWFAEL SVDMEKILEI IRVILKLYEQ
	WKNFDERKEM ATILSKMPKP KPPPNSEGEQ GPNGSQNSSY SQS Sequence without tag. The
	proposed Purification-Tag is based on experiences with the expression system, a different
	complexity of the protein could make another tag necessary. In case you have a special
	request, please contact us.
Characteristics:	Key Benefits:
	Made to order protein - from design to production - by highly experienced protein experts.

- · Protein expressed in mammalien cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

Grade:

custom-made

Target Details

Target:	Cyclin C (CCNC)
Alternative Name:	CCNC (CCNC Products)
Background:	Cyclin-C (SRB11 homolog) (hSRB11),FUNCTION: Component of the Mediator complex, a coactivator involved in regulated gene transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. Binds to and activates cyclin-dependent kinase CDK8 that phosphorylates the CTD (C-terminal domain) of the large subunit of RNA polymerase II (RNAp II), which may inhibit the formation of a transcription initiation complex. {ECO:0000269 PubMed:16595664, ECO:0000269 PubMed:8700522}.
Molecular Weight:	33.2 kDa
UniProt:	P24863
Pathways:	Cell Division Cycle, Regulation of Lipid Metabolism by PPARalpha

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months