# antibodies -online.com







# anti-BAG2 antibody (AA 151-211)





| $\sim$ |          |
|--------|----------|
| ( )\/\ | rview    |
| $\cup$ | 1 410 44 |

| edded |
|-------|
| ′     |

# **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human BAG2 |
|-----------------------|--|
| Isotype:              | IgG  |
| Cross-Reactivity:     | Human, Rat   |
| Predicted Reactivity: | Mouse,Dog,Cow,Pig,Horse                                  |
| Purification:         | Purified by Protein A.                                   |
|                       |  |

# **Target Details**

Target: BAG2


# Target Details

| Alternative Name:   | BAG2 (BAG2 Products)  |
|---------------------|---|
| Background:         | Synonyms: BAG-2, dJ417I1.2, BAG family molecular chaperone regulator 2, Bcl-2-associated athanogene 2, BAG2 |
|                     | Background: BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain                      |
|                     | and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG                 |
|                     | domain near the C terminus but differ markedly in their N-terminal regions. The predicted BAG2              |
|                     | protein contains 211 amino acids. The BAG domains of BAG1, BAG2, and BAG3 interact                          |
|                     | specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind              |
|                     | with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-               |
|                     | repressible manner. [provided by RefSeq].   |
| Gene ID:            | 9532  |
| UniProt:            | 095816  |
| Application Details |   |
| Application Notes:  | WB 1:300-5000   |
|                     | ELISA 1:500-1000  |
|                     | IHC-P 1:200-400   |
|                     | IHC-F 1:100-500   |
|                     | IF(IHC-P) 1:50-200  |
|                     | IF(IHC-F) 1:50-200  |
|                     | IF(ICC) 1:50-200  |
| Restrictions:       | For Research Use only   |
| Handling            |   |
| Format:             | Liquid  |
| Concentration:      | 1 μg/μL   |
| Buffer:             | 0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.                                       |
| Preservative:       | ProClin   |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be                         |
|                     | handled by trained staff only.  |
| Storage:            | 4 °C,-20 °C   |

# Handling

| Storage Comment: | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |
|------------------|---|
| Expiry Date:     | 12 months   |

# **Images**



# **Immunohistochemistry**

**Image 1.** Formalin-fixed and paraffin embedded rat brain labeled with Rabbit Anti BAG2 Polyclonal Antibody, Unconjugated (ABIN755347) at 1:200 followed by conjugation to the secondary antibody and DAB staining