

Datasheet for ABIN7553476

## NCAPD2 Protein (AA 1-1401) (His tag)



[Go to Product page](#)

### Overview

Quantity:	1 mg
Target:	NCAPD2
Protein Characteristics:	AA 1-1401
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This NCAPD2 protein is labelled with His tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB)

### Product Details

Purpose:	Custom-made recombinat NCAPD2 Protein expressed in mammalian cells.
Sequence:	<p>MAPQMYEFHL PLSPEELLKS GGVNQYVVQE VLSIKHLPPQ LRAFQAAFRA QGPLAMLQHF</p> <p>DTIYSILHHF RSIDPGLKED TLQFLIKVVS RHSQELPAIL DDTTLSGSDR NAHLNALKMN</p> <p>CYALIRLLES FETMASQTNL VDLDLGGKGK KARTKAAHGF DWEEERQPIL QLLTQLQLD</p> <p>IRHLWNHSII EEEFVSLVTG CCYRLLENPT INHQKNRPTR EAITHLLGVA LTRYNHMLSA</p> <p>TVKIIQMLQH FEHLAPVLVA AVSLWATDYG MKSIVGEIVR EIGQKCPQEL SRDPSGKGF</p> <p>AAFLTELAER VPAILMSSMC ILLDHLGGEN YMMRNAVLA MAEMVLQVLS GDQLEAAARD</p> <p>TRDQFLDTLQ AHGHDVNSFV RSRVLQLFTR IVQQKALPLT RFQAVVALAV GRLADKSVLV</p> <p>CKNAIQLLAS FLANNPFSCK LSDADLAGPL QKETQKLQEM RAQRRTAAAS AVLDPEEEWE</p> <p>AMLPELKSTL QLLQLPQGE EEIPEQIANT ETEDVKGRI YQLLAKASYK KAILTREAT</p> <p>GHFQESEPFS HIDPEESEET RLLNILGLIF KGPAASTQEK NPRESTGNMV TGQTVCKNKP</p> <p>NMSDPEESRG NDELVKQEML VQYLQDAYSF SRKITEAIGI ISKMMYENTT TVVQEVIEFF</p>

VMVFQFGVPQ ALFGVRRMLP LIWSKEPGVR EAVLNAYRQL YLNPKGDSAR AKAQALIQNL  
SLLLVDASVG TIQCLEEILC EFVQKDELKP AVTQLLWERA TEKVACCPLE RCSSVMLLGM  
MARGKPEIVG SNLDTLVSIG LDEKFPQDYR LAQQVCHAIA NISDRRKPSL GKRHPPFRLP  
QEHLRFERLR ETVTKGFVHP DPLWIPFKEV AVTLIYQLAE GPEVICAQIL QGCAKQALEK  
LEEKRTSQED PKESPAMLPT FLLMNLLSLA GDVALQQLVH LEQAVSGELC RRRVLREEQE  
HKTKDPKEKN TSSETTMEEE LGLVGATADD TEAELIRGIC EMELLDGKQT LAAFVPLLLK  
VCNNPGLYSN PDLSAAASLA LGKFCMISAT FCDSQLRLLF TMLEKSPLPI VRSNLMVATG  
DLAIRFPNLV DPWTPHLYAR LRDPAQQVRK TAGLVMTHLI LKDMVKVKGQ VSEMAVLLID  
PEPQIAALAK NFFNELSHKG NAIYNLLPDI ISRLSDPELG VEEEPFHTIM KQLLSYITKD  
KQTESLVEKL CQRFRTSRTE RQQRDLAYCV SQLPLTERGL RKMLDNFDCF GDKLSDSEIF  
SAFLSVVGKL RRGAKPEGKA IIDEFEQKLR ACHTRGLDGI KELEIGQAGS QRAPSAKKPS  
TGSRYQPLAS TASNDFVTP EPRRTTRRHP NTQQRASKKK PKVVFSSDES SEEDLSAEMT  
EETPKKTP ILRASARRHR S

**Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.**

Characteristics:

Key Benefits:

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

Grade:

custom-made

Target Details

Target:

NCAPD2

## Target Details

Alternative Name:	NCAPD2 ( <a href="#">NCAPD2 Products</a> )
Background:	<p>Condensin complex subunit 1 (Chromosome condensation-related SMC-associated protein 1) (Chromosome-associated protein D2) (hCAP-D2) (Non-SMC condensin I complex subunit D2) (XCAP-D2 homolog),FUNCTION: Regulatory subunit of the condensin complex, a complex required for conversion of interphase chromatin into mitotic-like condense chromosomes. The condensin complex probably introduces positive supercoils into relaxed DNA in the presence of type I topoisomerases and converts nicked DNA into positive knotted forms in the presence of type II topoisomerases. May target the condensin complex to DNA via its C-terminal domain (PubMed:11136719). May promote the resolution of double-strand DNA catenanes (intertwines) between sister chromatids. Condensin-mediated compaction likely increases tension in catenated sister chromatids, providing directionality for type II topoisomerase-mediated strand exchanges toward chromatid decatenation. Required for decatenation of non-centromeric ultrafine DNA bridges during anaphase. Early in neurogenesis, may play an essential role to ensure accurate mitotic chromosome condensation in neuron stem cells, ultimately affecting neuron pool and cortex size (PubMed:27737959).</p> <p>{ECO:0000269 PubMed:11136719, ECO:0000269 PubMed:27737959}.</p>
Molecular Weight:	157.2 kDa
UniProt:	<a href="#">Q15021</a>

## Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months