

Datasheet for ABIN7553747

DYNLT1 Protein (AA 1-113) (His tag)



Overview

Quantity:	1 mg
Target:	DYNLT1
Protein Characteristics:	AA 1-113
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This DYNLT1 protein is labelled with His tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB)
Product Details	
Purpose:	Custom-made recombinat DYNLT1 Protein expressed in mammalien cells.
Sequence:	MEDYQAAEET AFVVDEVSNI VKEAIESAIG GNAYQHSKVN QWTTNVVEQT LSQLTKLGKP
	FKYIVTCVIM QKNGAGLHTA SSCFWDSSTD GSCTVRWENK TMYCIVSAFG LSI Sequence
	without tag. The proposed Purification-Tag is based on experiences with the expression
	system, a different complexity of the protein could make another tag necessary. In case you
	have a special request, please contact us.
Characteristics:	Key Benefits:

• State-of-the-art algorithm used for plasmid design (Gene synthesis).

transmembrane proteins.

Made to order protein - from design to production - by highly experienced protein experts.
Protein expressed in mammalien cells and purified in one-step affinity chromatography
The optimized expression system ensures reliability for intracellular, secreted and

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

Grade:

custom-made

Target Details

Target: DYNLT1

Alternative Name: DYNLT1 (DYNLT1 Products)

Background:

homolog),FUNCTION: Acts as one of several non-catalytic accessory components of the cytoplasmic dynein 1 complex that are thought to be involved in linking dynein to cargos and to adapter proteins that regulate dynein function. Cytoplasmic dynein 1 acts as a motor for the intracellular retrograde motility of vesicles and organelles along microtubules. Binds to transport cargos and is involved in apical cargo transport such as rhodopsin-bearing vesicles in polarized epithelia. May also be a accessory component of axonemal dynein., FUNCTION: Plays a role in neuronal morphogenesis, the function is independent of cytoplasmic dynein and

seems to be coupled to regulation of the actin cytoskeleton by enhancing Rac1 activity. The

function in neurogenesis may be regulated by association with a G-protein beta-gamma dimer.

Dynein light chain Tctex-type 1 (Protein CW-1) (T-complex testis-specific protein 1

May function as a receptor-independent activator of heterotrimeric G-protein signaling, the activation appears to be independent of a nucleotide exchange. Plays a role in regulating neurogenesis, inhibits the genesis of neurons from precursor cells during cortical development presumably by antagonizing ARHGEF2. Involved in the regulation of mitotic spindle orientation (By similarity). Unrelated to the role in retrograde microtubule-associated movement may play a role in the dimerization of cytoplasmic proteins/domains such as for ACVR2B. Binds to the

{ECO:0000250, ECO:0000269|PubMed:27502274}., FUNCTION: (Microbial infection) Is involved

cytoplasmic domain of ACVR2B and, in vitro, inhibits ACVR2B signaling (PubMed:27502274).

in intracellular targeting of D-type retrovirus gag polyproteins to the cytoplasmic assembly site.

Target Details

rarget Details	
	{ECO:0000269 PubMed:18647839}.
Molecular Weight:	12.5 kDa
UniProt:	P63172
Pathways:	Regulation of G-Protein Coupled Receptor Protein Signaling
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a
	guarantee though.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months