

Datasheet for ABIN7554076
HDAC3 Protein (AA 1-428) (His tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	HDAC3
Protein Characteristics:	AA 1-428
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This HDAC3 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Purpose:	Custom-made recombinat HDAC3 Protein expressed in mammalian cells.
Sequence:	<p>MAKTVAYFYD PDVGNFHYGA GHMKPHRLA LTHSLVLHYG LYKMKIVFKP YQASQHDMCR FHSEYIDFL QRVSPNTMQG FTKSLNAFNV GDDCPVFPGL FEFCSRYTGA SLQGATQLNN KICDIANWA GGLHHAKKFE ASGFCYVNDI VIGILELLKY HPRVLYIDID IHHGDGVQEA FYLTDRVMTV SFHKYGNFYFF PGTGDMYEVG AESGRYYCLN VPLRDGIDDQ SYKHLFQPVI NQVDFYQPT CIVLQCGADS LGCDRLGCFN LSIRGHGECV EYVKSFNIP LVLGGGGYTV RNVARCWTYE TSLLVEEAIS EELPYSEYFE YFAPDFTLHP DVSTRIENQN SRQYLDQIRQ TIFENLKMLN HAPSVQIHDV PADLLTYDRT DEADAEERGP EENYSRPEAP NEFYDGDHDN DKESDVEI Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.</p>
Characteristics:	Key Benefits:

Product Details

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:	> 90 % as determined by Bis-Tris Page, Western Blot
---------	---

Grade:	custom-made
--------	-------------

Target Details

Target:	HDAC3
---------	-------

Alternative Name:	HDAC3 (HDAC3 Products)
-------------------	--

Background:	<p>Histone deacetylase 3 (HD3) (EC 3.5.1.98) (Protein deacetylase HDAC3) (EC 3.5.1.-) (Protein deacetylase HDAC3) (EC 3.5.1.-) (RPD3-2) (SMAP45),FUNCTION: Histone deacetylase that catalyzes the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4), and some other non-histone substrates (PubMed:23911289, PubMed:21030595, PubMed:21444723, PubMed:25301942, PubMed:28497810, PubMed:28167758, PubMed:32404892). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events (PubMed:23911289). Histone deacetylases act via the formation of large multiprotein complexes (PubMed:23911289). Participates in the BCL6 transcriptional repressor activity by deacetylating the H3 'Lys-27' (H3K27) on enhancer elements, antagonizing EP300 acetyltransferase activity and repressing proximal gene expression (PubMed:23911289). Acts as a molecular chaperone for shuttling phosphorylated NR2C1 to PML bodies for sumoylation (By similarity). Contributes, together with XBP1 isoform 1, to the activation of NFE2L2-mediated HMOX1 transcription factor gene expression in a PI(3)K/mTORC2/Akt-dependent signaling</p>
-------------	--

Target Details

pathway leading to endothelial cell (EC) survival under disturbed flow/oxidative stress (PubMed:25190803). Regulates both the transcriptional activation and repression phases of the circadian clock in a deacetylase activity-independent manner (By similarity). During the activation phase, promotes the accumulation of ubiquitinated BMAL1 at the E-boxes and during the repression phase, blocks FBXL3-mediated CRY1/2 ubiquitination and promotes the interaction of CRY1 and BMAL1 (By similarity). The NCOR1-HDAC3 complex regulates the circadian expression of the core clock gene BMAL1 and the genes involved in lipid metabolism in the liver (By similarity). Also functions as a deacetylase for non-histone targets, such as KAT5, MEF2D, MAPK14, RARA and STAT3 (PubMed:15653507, PubMed:21030595, PubMed:21444723, PubMed:25301942, PubMed:28167758). Serves as a corepressor of RARA, mediating its deacetylation and repression, leading to inhibition of RARE DNA element binding (PubMed:28167758). In association with RARA, plays a role in the repression of microRNA-10a and thereby in the inflammatory response (PubMed:28167758). In addition to protein deacetylase activity, also acts as a protein-lysine deacylase by recognizing other acyl groups: catalyzes removal of (2E)-butenoyl (crotonyl) and 2-hydroxyisobutanoyl (2-hydroxyisobutyryl) acyl groups from lysine residues, leading to protein decrotonylation and de-2-hydroxyisobutyrylation, respectively (PubMed:28497810, PubMed:29192674, PubMed:34608293). Catalyzes decrotonylation of MAPRE1/EB1 (PubMed:34608293). {ECO:0000250|UniProtKB:O88895, ECO:0000269|PubMed:15653507, ECO:0000269|PubMed:21030595, ECO:0000269|PubMed:21444723, ECO:0000269|PubMed:23911289, ECO:0000269|PubMed:25190803, ECO:0000269|PubMed:25301942, ECO:0000269|PubMed:28167758, ECO:0000269|PubMed:28497810, ECO:0000269|PubMed:29192674, ECO:0000269|PubMed:32404892, ECO:0000269|PubMed:34608293}.

Molecular Weight: 48.8 kDa

UniProt: [O15379](#)

Pathways: [Neurotrophin Signaling Pathway](#), [Regulation of Lipid Metabolism by PPARalpha](#), [Regulation of Muscle Cell Differentiation](#), [Skeletal Muscle Fiber Development](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months