

Datasheet for ABIN7554222

Importin 8 Protein (IPO8) (AA 1-1037) (His tag)



Overview

Quantity:	1 mg
Target:	Importin 8 (IPO8)
Protein Characteristics:	AA 1-1037
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Importin 8 protein is labelled with His tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB)

Product Details

Product Details	
Purpose:	Custom-made recombinat IPO8 Protein expressed in mammalien cells.
Sequence:	MDLNRIIQAL KGTIDPKLRI AAENELNQSY KIINFAPSLL RIIVSDHVEF PVRQAAAIYL
	KNMVTQYWPD REPPPGEAIF PFNIHENDRQ QIRDNIVEGI IRSPDLVRVQ LTMCLRAIIK
	HDFPGHWPGV VDKIDYYLQS QSSASWLGSL LCLYQLVKTY EYKKAEEREP LIIAMQIFLP
	RIQQQIVQLL PDSSYYSVLL QKQILKIFYA LVQYALPLQL VNNQTMTTWM EIFRTIIDRT
	VPPETLHIDE DDRPELVWWK CKKWALHIVA RLFERYGSPG NVTKEYFEFS EFFLKTYAVG
	IQQVLLKILD QYRQKEYVAP RVLQQAFNYL NQGVVHSITW KQMKPHIQNI SEDVIFSVMC
	YKDEDEELWQ EDPYEYIRMK FDIFEDYASP TTAAQTLLYT AAKKRKEVLP KMMAFCYQIL
	TDPNFDPRKK DGALHVIGSL AEILLKKSLF KDQMELFLQN HVFPLLLSNL GYLRARSCWV
	LHAFSSLKFH NELNLRNAVE LAKKSLIEDK EMPVKVEAAL ALQSLISNQI QAKEYMKPHV
	RPIMQELLHI VRETENDDVT NVIQKMICEY SQEVASIAVD MTQHLAEIFG KVLQSDEYEE
	VEDKTVMAMG ILHTIDTILT VVEDHKEITQ QLENICLRII DLVLQKHVIE FYEEILSLAY SLTCHSISPO

MWQLLGILYE VFQQDCFEYF TDMMPLLHNY VTIDTDTLLS NAKHLEILFT MCRKVLCGDA GEDAECHAAK LLEVIILQCK GRGIDQCIPL FVQLVLERLT RGVKTSELRT MCLQVAIAAL YYNPDLLLHT LERIQLPHNP GPITVQFINQ WMNDTDCFLG HHDRKMCIIG LSILLELQNR PPAVDAVVGQ IVPSILFLFL GLKQVCATRQ LVNREDRSKA EKADMEENEE ISSDEEETNV TAQAMQSNNG RGEDEEEDD DWDEEVLEET ALEGFSTPLD LDNSVDEYQF FTQALITVQS RDAAWYQLLM APLSEDQRTA LQEVYTLAEH RRTVAEAKKK IEQQGGFTFE NKGVLSAFNF GTVPSNN Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made to order protein from design to production by highly experienced protein experts.
- · Protein expressed in mammalien cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

Grade:

custom-made

Target Details

Target:	Importin 8 (IPO8)
Alternative Name:	IPO8 (IPO8 Products)
Background:	Importin-8 (Imp8) (Ran-binding protein 8) (RanBP8),FUNCTION: Involved in nuclear protein
	import, either by acting as autonomous nuclear transport receptor or as an adapter-like protein
	in association with the importin-beta subunit KPNB1. Acting autonomously, may serve as
	receptor for nuclear localization signals (NLS) and promote translocation of import substrates

through the nuclear pore complex (NPC) by an energy requiring, Ran-dependent mechanism. At the nucleoplasmic side of the NPC, Ran binds to importin, the importin/substrate complex dissociates and importin is re-exported from the nucleus to the cytoplasm where GTP hydrolysis releases Ran. The directionality of nuclear import is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus (PubMed:9214382). In vitro mediates the nuclear import of the signal recognition particle protein SRP19 (PubMed:11682607). May also be involved in cytoplasm-to-nucleus shuttling of a broad spectrum of other cargos, including Argonaute-microRNAs complexes, the JUN protein, RELA/NF-kappa-B p65 subunit, the translation initiation factor EIF4E and a set of receptor-activated mothers against decapentaplegic homolog (SMAD) transcription factors that play a critical role downstream of the large family of transforming growth factor beta and bone morphogenetic protein (BMP) cytokines (Probable). {ECO:0000269|PubMed:11682607, ECO:0000269|PubMed:9214382, ECO:0000305|PubMed:34010604}.

Molecular Weight:

119.9 kDa

UniProt:

015397

Application Details

Application Notes:

In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions:

For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months