

Datasheet for ABIN7554227

## JAK3 Protein (AA 1-1124) (His tag)



[Go to Product page](#)

### Overview

Quantity:	1 mg
Target:	JAK3
Protein Characteristics:	AA 1-1124
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This JAK3 protein is labelled with His tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB)

### Product Details

Purpose:	Custom-made recombinat JAK3 Protein expressed in mammalian cells.
Sequence:	<p>MAPPSEETPL IPQRSCSLLS TEAGALHVLL PARGPGPPQR LSFSFGDHLA EDLCVQAAKA</p> <p>SGILPVYHSL FALATEDLSC WFPPSHIFSV EDASTQVLLY RIRFYFPNWF GLEKCHRFGL</p> <p>RKDLASAILD LPVLEHLFAQ HRSDLVSGRL PVGLSLKEQG ECLSLAVLDL ARMAREQAQR</p> <p>PGELLKTVSY KACLPPLSRD LIQGLSFVTR RRIRRTVRR LRRVAACQAD RHSLMAKYIM</p> <p>DLERLDPAGA AETFHVGLPG ALGGHDGLGL LRVAGDGGIA WTQGEQEV LQ PFCDFPEIVD</p> <p>ISIKQAPRVG PAGEHRLVTV TRTDNQILEA EFPGLPEALS FVALVDGYFR LTDSQHFFC</p> <p>KEVAPRLL E EVAEQCHGPI TLDFAINKLK TGGSRPGSYV LRRSPQDFDS FLLTVCVQNP</p> <p>LGPDYKGCLI RRSPTGTFL VGLSRPHSSL RELLATCWDG GLHVDGVA VT LTSCCIPRPK</p> <p>EKSNLIVVQR GHSPPTSSLV QPQSQYQLSQ MTFHKIPADS LEWHENLG HG SFTKIYRGCR</p> <p>HEVVDGEARK TEVLLKVMDA KHKNCMESFL EAASLMSQVS YRHLVLLHGV CMAGDSTMVQ</p> <p>EFVHLGAIDM YLRKRGHLVP ASWKLQVVKQ LAYALNYLED KGLPHGNVSA RKVLLAREGA</p>

DGSPFFIKLS DPGVSPAVLS LEMLTDRIPW VAPECLREAQ TLSLEADKWG FGATVWEVFS  
GVTMPISALD PAKKLQFYED RQQLPAPKWT ELALLIQQCM AYEPVQRPSF RAVIRDLNSL  
ISSDYELLSD PTPGALAPRD GLWNGAQLYA CQDPTIFEER HLKYISQLGK GNFGSVELCR  
YDPLGDNTGA LVAVKQLQHS GPDQQRDFQR EIQLKALHS DFIVKYRGVS YGPGRQSLRL  
VMEYLPSCGL RDFLQRHRAR LDASRLLLYS SQICKGMEYL GSRRCVHRDL AARNILVESE  
AHVKIADFGL AKLLPLDKDY YVVREPGQSP IFWYAPESLS DNIFSRQSDV WSFGVVLYEL  
FTYCDKSCSP SAEFLRMMGC ERDVPALCRL LELLEEGQRL PAPPACPAEV HELMKLCWAP  
SPQDRPSFSA LGPQLDMLWS GSRGCETHAF TAHPEGKHHS LSFS **Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.**

Characteristics:	<p>Key Benefits:</p> <ul style="list-style-type: none"><li>• Made to order protein - from design to production - by highly experienced protein experts.</li><li>• Protein expressed in mammalian cells and purified in one-step affinity chromatography</li><li>• The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.</li><li>• State-of-the-art algorithm used for plasmid design (Gene synthesis).</li></ul> <p>This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.</p> <p>If you are not interested in a full length protein, please contact us for individual protein fragments.</p> <p>The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.</p>
------------------	---

Purity:	> 90 % as determined by Bis-Tris Page, Western Blot
Grade:	custom-made

Target Details

Target:	JAK3
Alternative Name:	JAK3 (JAK3 Products)
Background:	Tyrosine-protein kinase JAK3 (EC 2.7.10.2) (Janus kinase 3) (JAK-3) (Leukocyte janus kinase) (L-JAK),FUNCTION: Non-receptor tyrosine kinase involved in various processes such as cell

## Target Details

growth, development, or differentiation. Mediates essential signaling events in both innate and adaptive immunity and plays a crucial role in hematopoiesis during T-cells development. In the cytoplasm, plays a pivotal role in signal transduction via its association with type I receptors sharing the common subunit gamma such as IL2R, IL4R, IL7R, IL9R, IL15R and IL21R. Following ligand binding to cell surface receptors, phosphorylates specific tyrosine residues on the cytoplasmic tails of the receptor, creating docking sites for STATs proteins. Subsequently, phosphorylates the STATs proteins once they are recruited to the receptor. Phosphorylated STATs then form homodimer or heterodimers and translocate to the nucleus to activate gene transcription. For example, upon IL2R activation by IL2, JAK1 and JAK3 Molecules bind to IL2R beta (IL2RB) and gamma chain (IL2RG) subunits inducing the tyrosine phosphorylation of both receptor subunits on their cytoplasmic domain. Then, STAT5A and STAT5B are recruited, phosphorylated and activated by JAK1 and JAK3. Once activated, dimerized STAT5 translocates to the nucleus and promotes the transcription of specific target genes in a cytokine-specific fashion. {ECO:0000269|PubMed:11909529, ECO:0000269|PubMed:20440074, ECO:0000269|PubMed:7662955, ECO:0000269|PubMed:8022485}.

Molecular Weight:	125.1 kDa
UniProt:	<a href="#">P52333</a>
Pathways:	<a href="#">JAK-STAT Signaling</a> , <a href="#">RTK Signaling</a> , <a href="#">Response to Growth Hormone Stimulus</a> , <a href="#">Regulation of Leukocyte Mediated Immunity</a> , <a href="#">Production of Molecular Mediator of Immune Response</a> , <a href="#">Protein targeting to Nucleus</a> , <a href="#">Activated T Cell Proliferation</a> , <a href="#">Unfolded Protein Response</a>

## Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C

Handling

---

Storage Comment: Store at -80°C.

---

Expiry Date: 12 months