

Datasheet for ABIN7554368 **KMO Protein (AA 1-486) (His tag)**



Overview

Quantity:	1 mg
Target:	KMO
Protein Characteristics:	AA 1-486
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This KMO protein is labelled with His tag.

Purpose:	Custom-made recombinant KMO Protein expressed in mammalian cells.
Sequence:	MDSSVIQRKK VAVIGGGLVG SLQACFLAKR NFQIDVYEAR EDTRVATFTR GRSINLALSH
	RGRQALKAVG LEDQIVSQGI PMRARMIHSL SGKKSAIPYG TKSQYILSVS RENLNKDLLT
	AAEKYPNVKM HFNHRLLKCN PEEGMITVLG SDKVPKDVTC DLIVGCDGAY STVRSHLMKK
	PRFDYSQQYI PHGYMELTIP PKNGDYAMEP NYLHIWPRNT FMMIALPNMN KSFTCTLFMP
	FEEFEKLLTS NDVVDFFQKY FPDAIPLIGE KLLVQDFFLL PAQPMISVKC SSFHFKSHCV
	LLGDAAHAIV PFFGQGMNAG FEDCLVFDEL MDKFSNDLSL CLPVFSRLRI PDDHAISDLS
	MYNYIEMRAH VNSSWFIFQK NMERFLHAIM PSTFIPLYTM VTFSRIRYHE AVQRWHWQKK
	VINKGLFFLG SLIAISSTYL LIHYMSPRSF LRLRRPWNWI AHFRNTTCFP AKAVDSLEQI SNLISR
	Sequence without tag. The proposed Purification-Tag is based on experiences with the
	expression system, a different complexity of the protein could make another tag necessary.
	In case you have a special request, please contact us.

Product Details

	isoform, please contact us regarding an individual offer.
Characteristics:	Key Benefits:
	Made to order protein - from design to production - by highly experienced protein experts.
	Protein expressed in mammalian cells and purified in one-step affinity chromatography
	The optimized expression system ensures reliability for intracellular, secreted and
	transmembrane proteins. • State-of-the-art algorithm used for plasmid design (Gene synthesis).
	State of the art algorithm asea for plasmia design (defic synthesis).
	This protein is a made-to-order protein and will be made for the first time for your order. Our
	experts in the lab try to ensure that you receive soluble protein.
	If you are not interested in a full length protein, please contact us for individual protein
	fragments.
	The big advantage of ordering our made-to-order proteins in comparison to ordering custom
	made proteins from other companies is that there is no financial obligation in case the protein
	cannot be expressed or purified.
Purity:	> 90 % as determined by Bis-Tris PAGE, anti-tag ELISA, Western Blot and analytical SEC (HPLC
Grade:	custom-made
Target Details	
Target:	
rarget.	KMO
	KMO (KMO Products)
Alternative Name:	
Alternative Name:	KMO (KMO Products)
Alternative Name:	KMO (KMO Products) Kynurenine 3-monooxygenase (EC 1.14.13.9) (Kynurenine 3-hydroxylase),FUNCTION: Catalyze
Alternative Name:	KMO (KMO Products) Kynurenine 3-monooxygenase (EC 1.14.13.9) (Kynurenine 3-hydroxylase),FUNCTION: Catalyze the hydroxylation of L-kynurenine (L-Kyn) to form 3-hydroxy-L-kynurenine (L-30HKyn)
Alternative Name:	KMO (KMO Products) Kynurenine 3-monooxygenase (EC 1.14.13.9) (Kynurenine 3-hydroxylase),FUNCTION: Catalyze the hydroxylation of L-kynurenine (L-Kyn) to form 3-hydroxy-L-kynurenine (L-30HKyn) (PubMed:29429898, PubMed:23575632, PubMed:26752518, PubMed:28604669,
Alternative Name:	KMO (KMO Products) Kynurenine 3-monooxygenase (EC 1.14.13.9) (Kynurenine 3-hydroxylase),FUNCTION: Catalyze the hydroxylation of L-kynurenine (L-Kyn) to form 3-hydroxy-L-kynurenine (L-30HKyn) (PubMed:29429898, PubMed:23575632, PubMed:26752518, PubMed:28604669, PubMed:29208702). Required for synthesis of quinolinic acid, a neurotoxic NMDA receptor
Alternative Name:	KMO (KMO Products) Kynurenine 3-monooxygenase (EC 1.14.13.9) (Kynurenine 3-hydroxylase),FUNCTION: Catalyze the hydroxylation of L-kynurenine (L-Kyn) to form 3-hydroxy-L-kynurenine (L-30HKyn) (PubMed:29429898, PubMed:23575632, PubMed:26752518, PubMed:28604669, PubMed:29208702). Required for synthesis of quinolinic acid, a neurotoxic NMDA receptor antagonist and potential endogenous inhibitor of NMDA receptor signaling in axonal targeting,
Alternative Name:	Kynurenine 3-monooxygenase (EC 1.14.13.9) (Kynurenine 3-hydroxylase),FUNCTION: Catalyze the hydroxylation of L-kynurenine (L-Kyn) to form 3-hydroxy-L-kynurenine (L-30HKyn) (PubMed:29429898, PubMed:23575632, PubMed:26752518, PubMed:28604669, PubMed:29208702). Required for synthesis of quinolinic acid, a neurotoxic NMDA receptor antagonist and potential endogenous inhibitor of NMDA receptor signaling in axonal targeting, synaptogenesis and apoptosis during brain development. Quinolinic acid may also affect
	Kynurenine 3-monooxygenase (EC 1.14.13.9) (Kynurenine 3-hydroxylase),FUNCTION: Catalyze the hydroxylation of L-kynurenine (L-Kyn) to form 3-hydroxy-L-kynurenine (L-30HKyn) (PubMed:29429898, PubMed:23575632, PubMed:26752518, PubMed:28604669, PubMed:29208702). Required for synthesis of quinolinic acid, a neurotoxic NMDA receptor antagonist and potential endogenous inhibitor of NMDA receptor signaling in axonal targeting, synaptogenesis and apoptosis during brain development. Quinolinic acid may also affect NMDA receptor signaling in pancreatic beta cells, osteoblasts, myocardial cells, and the
Alternative Name:	Kynurenine 3-monooxygenase (EC 1.14.13.9) (Kynurenine 3-hydroxylase),FUNCTION: Catalyze the hydroxylation of L-kynurenine (L-Kyn) to form 3-hydroxy-L-kynurenine (L-30HKyn) (PubMed:29429898, PubMed:23575632, PubMed:26752518, PubMed:28604669, PubMed:29208702). Required for synthesis of quinolinic acid, a neurotoxic NMDA receptor antagonist and potential endogenous inhibitor of NMDA receptor signaling in axonal targeting, synaptogenesis and apoptosis during brain development. Quinolinic acid may also affect NMDA receptor signaling in pancreatic beta cells, osteoblasts, myocardial cells, and the gastrointestinal tract (Probable). {ECO:0000269 PubMed:23575632,
Alternative Name:	Kynurenine 3-monooxygenase (EC 1.14.13.9) (Kynurenine 3-hydroxylase),FUNCTION: Catalyze the hydroxylation of L-kynurenine (L-Kyn) to form 3-hydroxy-L-kynurenine (L-30HKyn) (PubMed:29429898, PubMed:23575632, PubMed:26752518, PubMed:28604669, PubMed:29208702). Required for synthesis of quinolinic acid, a neurotoxic NMDA receptor antagonist and potential endogenous inhibitor of NMDA receptor signaling in axonal targeting, synaptogenesis and apoptosis during brain development. Quinolinic acid may also affect NMDA receptor signaling in pancreatic beta cells, osteoblasts, myocardial cells, and the gastrointestinal tract (Probable). {ECO:0000269 PubMed:23575632, ECO:0000269 PubMed:26752518, ECO:0000269 PubMed:28604669,

Target Details UniProt: 015229 **Application Details** We expect the protein to work for functional studies. As the protein has not been tested for Application Notes: functional studies yet we cannot offer a guarantee though. Restrictions: For Research Use only Handling Format: Liquid Buffer: The buffer composition is at the discretion of the manufacturer. Avoid repeated freeze-thaw cycles. Handling Advice:

-80 °C

Store at -80°C.

12 months

Storage:

Expiry Date:

Storage Comment: