

Datasheet for ABIN7554570 MAGOH Protein (AA 1-146) (His tag)



('	١١/		rv		۱۸/

Quantity:	1 mg
Target:	MAGOH
Protein Characteristics:	AA 1-146
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This MAGOH protein is labelled with His tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB)
Product Details	
Purpose:	Custom-made recombinat MAGOH Protein expressed in mammalien cells.
Sequence:	MESDFYLRYY VGHKGKFGHE FLEFEFRPDG KLRYANNSNY KNDVMIRKEA YVHKSVMEEL
	KRIIDDSEIT KEDDALWPPP DRVGRQELEI VIGDEHISFT TSKIGSLIDV NQSKDPEGLR
	VFYYLVQDLK CLVFSLIGLH FKIKPI Sequence without tag. The proposed Purification-Tag is
	based on experiences with the expression system, a different complexity of the protein
	could make another tag necessary. In case you have a special request, please contact us.
Characteristics:	Key Benefits:
	Made to order protein - from design to production - by highly experienced protein experts.
	Protein expressed in mammalien cells and purified in one-step affinity chromatography
	 The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
	State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

Grade:

custom-made

Target Details

Target:	MAGOH
. u. got.	

Alternative Name:

MAGOH (MAGOH Products)

Background:

Protein mago nashi homolog, FUNCTION: Required for pre-mRNA splicing as component of the spliceosome (PubMed:11991638). Plays a redundant role with MAGOHB as core component of the exon junction complex (EJC) and in the nonsense-mediated decay (NMD) pathway (PubMed:23917022). The EJC is a dynamic structure consisting of core proteins and several peripheral nuclear and cytoplasmic associated factors that join the complex only transiently either during EJC assembly or during subsequent mRNA metabolism. The EJC marks the position of the exon-exon junction in the mature mRNA for the gene expression machinery and the core components remain bound to spliced mRNAs throughout all stages of mRNA metabolism thereby influencing downstream processes including nuclear mRNA export, subcellular mRNA localization, translation efficiency and nonsense-mediated mRNA decay (NMD). The MAGOH-RBM8A heterodimer inhibits the ATPase activity of EIF4A3, thereby trapping the ATP-bound EJC core onto spliced mRNA in a stable conformation. The MAGOH-RBM8A heterodimer interacts with the EJC key regulator PYM1 leading to EJC disassembly in the cytoplasm and translation enhancement of EJC-bearing spliced mRNAs by recruiting them to the ribosomal 48S preinitiation complex. Involved in the splicing modulation of BCL2L1/Bcl-X (and probably other apoptotic genes), specifically inhibits formation of proapoptotic isoforms such as Bcl-X(S), the function is different from the established EJC assembly. {ECO:0000269|PubMed:11991638, ECO:0000269|PubMed:12730685, ECO:0000269|PubMed:16209946, ECO:0000269|PubMed:22203037,

Target Details

Expiry Date:

12 months

	ECO:0000269 PubMed:23917022}.
Molecular Weight:	17.2 kDa
UniProt:	P61326
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.