

Datasheet for ABIN7554805 NR4A3 Protein (AA 1-626) (His tag)



Overview

Quantity:	1 mg
Target:	NR4A3
Protein Characteristics:	AA 1-626
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This NR4A3 protein is labelled with His tag.

Product Details

Purpose:	Custom-made recombinant NR4A3 Protein expressed in mammalian cells.
Sequence:	MPCVQAQYSP SPPGSSYAAQ TYSSEYTTEI MNPDYTKLTM DLGSTEITAT ATTSLPSIST
	FVEGYSSNYE LKPSCVYQMQ RPLIKVEEGR APSYHHHHHH HHHHHHHHQQ QHQQPSIPPA
	SSPEDEVLPS TSMYFKQSPP STPTTPAFPP QAGALWDEAL PSAPGCIAPG PLLDPPMKAV
	PTVAGARFPL FHFKPSPPHP PAPSPAGGHH LGYDPTAAAA LSLPLGAAAA AGSQAAALES
	HPYGLPLAKR AAPLAFPPLG LTPSPTASSL LGESPSLPSP PSRSSSSGEG TCAVCGDNAA
	CQHYGVRTCE GCKGFFKRTV QKNAKYVCLA NKNCPVDKRR RNRCQYCRFQ KCLSVGMVKE
	VVRTDSLKGR RGRLPSKPKS PLQQEPSQPS PPSPPICMMN ALVRALTDST PRDLDYSRYC
	PTDQAAAGTD AEHVQQFYNL LTASIDVSRS WAEKIPGFTD LPKEDQTLLI ESAFLELFVL
	RLSIRSNTAE DKFVFCNGLV LHRLQCLRGF GEWLDSIKDF SLNLQSLNLD IQALACLSAL
	SMITERHGLK EPKRVEELCN KITSSLKDHQ SKGQALEPTE SKVLGALVEL RKICTLGLQR
	IFYLKLEDLV SPPSIIDKLF LDTLPF Sequence without tag. The proposed Purification-Tag is
	based on experiences with the expression system, a different complexity of the protein

	could make another tag necessary. In case you have a special request, please contact us.
Specificity:	If you are looking for a specific domain and are interested in a partial protein or a different
	isoform, please contact us regarding an individual offer.
Characteristics:	Key Benefits:
	 Made to order protein - from design to production - by highly experienced protein experts. Protein expressed in mammalian cells and purified in one-step affinity chromatography The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
	State-of-the-art algorithm used for plasmid design (Gene synthesis).
	This protein is a made-to-order protein and will be made for the first time for your order. Our
	experts in the lab try to ensure that you receive soluble protein.
	If you are not interested in a full length protein, please contact us for individual protein
	fragments.
	The big advantage of ordering our made-to-order proteins in comparison to ordering custom
	made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.
Purity:	> 90 % as determined by Bis-Tris PAGE, anti-tag ELISA, Western Blot and analytical SEC (HPLC
Grade:	custom-made
Target Details	
Farget:	NR4A3
Alternative Name:	NR4A3 (NR4A3 Products)
Background:	Nuclear receptor subfamily 4 group A member 3 (Mitogen-induced nuclear orphan receptor)
	(Neuron-derived orphan receptor 1) (Nuclear hormone receptor NOR-1),FUNCTION:
	Transcriptional activator that binds to regulatory elements in promoter regions in a cell- and
	response element (target)-specific manner. Induces gene expression by binding as monomer
	to the NR4A1 response element (NBRE) 5'-AAAAGGTCA-3' site and as homodimers to the Nu

muscle, myeloid progenitor cell and type B pancreatic cells, promotes mitogen-induced

vascular smooth muscle cell proliferation through transactivation of SKP2 promoter by binding

a NBRE site (By similarity). Upon PDGF stimulation, stimulates vascular smooth muscle cell proliferation by regulating CCND1 and CCND2 expression. In islets, induces type B pancreatic cell proliferation through up-regulation of genes that activate cell cycle, as well as genes that cause degradation of the CDKN1A (By similarity). Negatively regulates myeloid progenitor cell proliferation by repressing RUNX1 in a NBRE site-independent manner. During inner ear, plays a role as a key mediator of the proliferative growth phase of semicircular canal development (By similarity). Mediates also survival of neuron and smooth muscle cells, mediates CREB-induced neuronal survival, and during hippocampus development, plays a critical role in pyramidal cell survival and axonal guidance. Is required for S phase entry of the cell cycle and survival of smooth muscle cells by inducing CCND1, resulting in RB1 phosphorylation. Binds to NBRE motif in CCND1 promoter, resulting in the activation of the promoter and CCND1 transcription (By similarity). Also plays a role in inflammation, upon TNF stimulation, mediates monocyte adhesion by inducing the expression of VCAM1 and ICAM1 by binding to the NBRE consensus site (By similarity) (PubMed:20558821). In mast cells activated by Fc-epsilon receptor crosslinking, promotes the synthesis and release of cytokines but impairs events leading to degranulation (By similarity). Also plays a role in metabolism, by modulating feeding behavior, and by playing a role in energy balance by inhibiting the glucocorticoid-induced orexigenic neuropeptides AGRP expression, at least in part by forming a complex with activated NR3C1 on the AGRP- glucocorticoid response element (GRE), and thus weakening the DNA binding activity of NR3C1. Upon catecholamines stimulation, regulates gene expression that controls oxidative metabolism in skeletal muscle (By similarity). Plays a role in glucose transport by regulating translocation of the SLC2A4 glucose transporter to the cell surface (PubMed:24022864). Finally, during gastrulation plays a crucial role in the formation of anterior mesoderm by controlling cell migration. Inhibits adipogenesis (By similarity). Also participates in cardiac hypertrophy by activating PARP1 (By similarity). (ECO:0000250|UniProtKB:P51179, ECO:0000250|UniProtKB:Q9QZB6, ECO:0000269|PubMed:20558821, ECO:0000269|PubMed:24022864}.

Molecular Weight:

68.2 kDa

UniProt:

Q92570

Pathways:

Fc-epsilon Receptor Signaling Pathway, Nuclear Receptor Transcription Pathway, Steroid Hormone Mediated Signaling Pathway

Application Details

Application Notes:

We expect the protein to work for functional studies. As the protein has not been tested for

Application Details

	functional studies yet we cannot offer a guarantee though.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months