

Datasheet for ABIN7554934  
**PER1 Protein (AA 1-1290) (His tag)**



[Go to Product page](#)

## Overview

Quantity:	1 mg
Target:	PER1
Protein Characteristics:	AA 1-1290
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This PER1 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

## Product Details

Purpose:	Custom-made recombinat PER1 Protein expressed in mammalien cells.
Sequence:	<p>MSGPLEGADG GGDPRPGESF CPGGVPSGP PQRHPCGPS LADTDANSN GSSGNESNGH  ESRGASQRSS HSSSSGNGKD SALLETTESS KSTNSQSPSP PSSSIAYSL SASSQDNPS  TSGCSSEQSA RARTQKELMT ALRELKLRP PERRGKGRSG TLATLQYALA CVKQVQANQE  YYQWSLEEG EPCSMDMSTY TLEELHITS EYTLQNQDTF SVAVSFLTGR IVYISEQAAV  LLRCKRDVFR GTRFSELLAP QDVGVFYGST APSRLPTWGT GASAGSGLRD FTQEKSVFCR  IRGGPDRDPG PRYQPFRLTP YVTKIRVSDG APAQPCLLI AERIHSYGIEA PRIPPKRIF  TTRHTPSCLF QDVDERAAPL LGYLPQDLLG APVLLFLHPE DRPLMLAIHK KILQLAGQPF  DHSPIRFCAR NGEYVTMDTS WAGFVHPWSR KVAFVLGRHK VRTAPLNEDV FTTPPASPAP  SLDTDIQELS EQIHRLLLQP VHSPSPTGLC GVGAVTSPGP LHSPGSSSDS NGGDAEGPGP  PAPVTFQIC KDVHLVKHQG QQLFIESRAR PQSRPRLPAT GTFKAKALPC QSPDPELEAG  SAPVQAPLAL VPEEAERKEA SSSCYQQINC LDSILRYLES CNLPSTTKRK CASSSSYTTS</p>

## Product Details

---

SASDDDRQRT GPVSVGTKKD PPSAALSSEG ATPRKEPVVG GTLSPLALAN KAESVVSPTS  
QCSFSSTIVH VGDKKPPESD IIMMEDLPGL APGPAPSPAP SPTVAPDPAP DAYRPVGLTK  
AVLSLHTQKE EQAFLSRFRD LGRLRGLDSS STAPSALGER GCHHGPPAPS RRHHCRSKAK  
RSRHHQNPR A EAPCYVSHPS PVPPSTPWPT PPATTPFAV VQPYPLPVFS PRGGPQPLPP  
APTSVPPAAF PAVLVTPMVA LVLPNYLFPT PSSYPYGALQ TPAEGPPTPA SHSPSPSLPA  
LAPSPPHRPD SPLFNSRCSS PLQLNLLQLE ELPRAEGAAV AGGPGSSAGP PPPSAEAAEP  
EARLAEVTES SNQDALSGSS DLLELLLQED SRSRGTSAA SGLGSLGSG SGSGSHEGGS  
TSASITRSSQ SSHTSKYFGS IDSSEAEAGA ARGGAEPGDQ VIKYVLQDPI WLLMANADQR  
VMMTYQVPSR DMTSVLKQDR ERLRAMQKQQ PRFSEDQRRE LGAVHSWVRK GQLPRALDVM  
ACVDCGSSTQ DPGHPDDPLF SELDGLGLEP MEEGGGEQGS SGGGSGELEG CEEAQGGAKA  
SSSQDLAMEE EEEGRSSSSP ALPTAGNCTS **Sequence without tag. The proposed Purification-  
Tag is based on experiences with the expression system, a different complexity of the  
protein could make another tag necessary. In case you have a special request, please  
contact us.**

---

### Characteristics:

#### Key Benefits:

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

---

### Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

---

### Grade:

custom-made

---

## Target Details

---

### Target:

PER1

---

## Target Details

---

Alternative Name: PER1 ([PER1 Products](#))

---

**Background:** Period circadian protein homolog 1 (hPER1) (Circadian clock protein PERIOD 1) (Circadian pacemaker protein Rigui),FUNCTION: Transcriptional repressor which forms a core component of the circadian clock. The circadian clock, an internal time-keeping system, regulates various physiological processes through the generation of approximately 24 hour circadian rhythms in gene expression, which are translated into rhythms in metabolism and behavior. It is derived from the Latin roots 'circa' (about) and 'diem' (day) and acts as an important regulator of a wide array of physiological functions including metabolism, sleep, body temperature, blood pressure, endocrine, immune, cardiovascular, and renal function. Consists of two major components: the central clock, residing in the suprachiasmatic nucleus (SCN) of the brain, and the peripheral clocks that are present in nearly every tissue and organ system. Both the central and peripheral clocks can be reset by environmental cues, also known as Zeitgebers (German for 'timegivers'). The predominant Zeitgeber for the central clock is light, which is sensed by retina and signals directly to the SCN. The central clock entrains the peripheral clocks through neuronal and hormonal signals, body temperature and feeding-related cues, aligning all clocks with the external light/dark cycle. Circadian rhythms allow an organism to achieve temporal homeostasis with its environment at the molecular level by regulating gene expression to create a peak of protein expression once every 24 hours to control when a particular physiological process is most active with respect to the solar day. Transcription and translation of core clock components (CLOCK, NPAS2, BMAL1, BMAL2, PER1, PER2, PER3, CRY1 and CRY2) plays a critical role in rhythm generation, whereas delays imposed by post-translational modifications (PTMs) are important for determining the period ( $\tau$ ) of the rhythms ( $\tau$  refers to the period of a rhythm and is the length, in time, of one complete cycle). A diurnal rhythm is synchronized with the day/night cycle, while the ultradian and infradian rhythms have a period shorter and longer than 24 hours, respectively. Disruptions in the circadian rhythms contribute to the pathology of cardiovascular diseases, cancer, metabolic syndromes and aging. A transcription/translation feedback loop (TTFL) forms the core of the molecular circadian clock mechanism. Transcription factors, CLOCK or NPAS2 and BMAL1 or BMAL2, form the positive limb of the feedback loop, act in the form of a heterodimer and activate the transcription of core clock genes and clock-controlled genes (involved in key metabolic processes), harboring E-box elements (5'-CACGTG-3') within their promoters. The core clock genes: PER1/2/3 and CRY1/2 which are transcriptional repressors form the negative limb of the feedback loop and interact with the CLOCK|NPAS2-BMAL1|BMAL2 heterodimer inhibiting its activity and thereby negatively regulating their own expression. This heterodimer also activates nuclear receptors NR1D1/2 and RORA/B/G, which form a second feedback loop and which activate and repress BMAL1

---

## Target Details

---

transcription, respectively. Regulates circadian target genes expression at post-transcriptional levels, but may not be required for the repression at transcriptional level. Controls PER2 protein decay. Represses CRY2 preventing its repression on CLOCK/BMAL1 target genes such as FXYD5 and SCNN1A in kidney and PPARA in liver. Besides its involvement in the maintenance of the circadian clock, has an important function in the regulation of several processes. Participates in the repression of glucocorticoid receptor NR3C1/GR-induced transcriptional activity by reducing the association of NR3C1/GR to glucocorticoid response elements (GREs) by BMAL1:CLOCK. Plays a role in the modulation of the neuroinflammatory state via the regulation of inflammatory mediators release, such as CCL2 and IL6. In spinal astrocytes, negatively regulates the MAPK14/p38 and MAPK8/JNK MAPK cascades as well as the subsequent activation of NFkappaB. Coordinately regulates the expression of multiple genes that are involved in the regulation of renal sodium reabsorption. Can act as gene expression activator in a gene and tissue specific manner, in kidney enhances WNK1 and SLC12A3 expression in collaboration with CLOCK. Modulates hair follicle cycling. Represses the CLOCK-BMAL1 induced transcription of BHLHE40/DEC1. {ECO:0000269|PubMed:24005054}.

---

Molecular Weight: 136.2 kDa

---

UniProt: [O15534](#)

---

Pathways: [Photoperiodism](#)

## Application Details

---

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

---

Buffer: The buffer composition is at the discretion of the manufacturer.

---

Handling Advice: Avoid repeated freeze-thaw cycles.

---

Storage: -80 °C

---

Storage Comment: Store at -80°C.

---

Expiry Date: 12 months