

Datasheet for ABIN7554976

PIWIL1 Protein (AA 1-861) (His tag)



[Go to Product page](#)

Overview

| | |
|-------------------------------|---|
| Quantity: | 1 mg |
| Target: | PIWIL1 |
| Protein Characteristics: | AA 1-861 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This PIWIL1 protein is labelled with His tag. |
| Application: | Western Blotting (WB), SDS-PAGE (SDS) |

Product Details

| | |
|-----------|--|
| Purpose: | Custom-made recombinat PIWIL1 Protein expressed in mammalian cells. |
| Sequence: | <p>MTGRARARAR GRARGQETAQ LVGSTASQQP GYIQRPPQPP PAEGELFGRG RQRGTAGGTA</p> <p>KSQGLQISAG FQELSLAERG GRRRDFHDLG VNTRQNLDHV KESKTGSSGI IVRLSTNHFR</p> <p>LTSRPQWALY QYHIDYNPLM EARRLSALL FQHEDLIGKC HAFDGTILFL PKRLQQKVTE</p> <p>VFSKTRNGED VRITITLTNE LPPTSPTCLQ FYNIIFRRL KIMNLQQIGR NYYNPNPID</p> <p>IPSHRLVIWP GFTTSILQYE NSIMLCTDVS HKVLRSETVL DFMFNFYHQT EEHKFQEQVS</p> <p>KELIGLVVLT KYNNKTYRVD DIDWDQNP KS TFKKADGSEV SFLEYRKYQY NQEITDLKQP</p> <p>VLVSQPKRRR GPGGTLPGPA MLIPELCYLT GLTDKMRNDF NVMKDLAVHT RLTPEQRQRE</p> <p>VGRLIDYIHK NDNVQREL RD WGLSFD SNLL SFSGRILQTE KIHQGGKTFD YNPQFADWSK</p> <p>ETRGAPLISV KPLDNWLLIY TRRNYEAANS LIQNLFKVTP AMGMQMRKAI MIEVDDRTEA</p> <p>YLRVLQQKV T ADTQIVVCLL SSNRKDKYDA IKKYLCTDCP TPSQCVVART LGKQQTVMAI</p> <p>ATKIALQMNC KMGGELWRVD IPLKLV MIVG IDCYHDMTAG RRSIAGFVAS INEGMTRWFS</p> |

Product Details

RCIFQDRGQE LVDGLKVCLQ AALRAWNSCN EYMPSRIIVY RDGVGDGQLK TLVNYEVPQF
LDCLKSIGRG YNPRLTVIVV KKRVNTRFFA QSGGRLQNPL PGTVIDVEVT RPEWYDFFIV
SQA VRSGSVS PTHYNVIYDN SGLKPDHIQR LTYKLCHIIY NWPGVIRVPA PCQYAHKLAF
LVGQSIHREP NLSLSNRLYY L

Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

Grade:

custom-made

Target Details

Target:

PIWIL1

Alternative Name:

PIWIL1 ([PIWIL1 Products](#))

Background:

Piwi-like protein 1 (EC 3.1.26.-),FUNCTION: Endoribonuclease that plays a central role in postnatal germ cells by repressing transposable elements and preventing their mobilization, which is essential for the germline integrity. Acts via the piRNA metabolic process, which mediates the repression of transposable elements during meiosis by forming complexes composed of piRNAs and Piwi proteins and governs the methylation and subsequent repression of transposons. Directly binds methylated piRNAs, a class of 24 to 30 nucleotide RNAs that are generated by a Dicer-independent mechanism and are primarily derived from

Target Details

transposons and other repeated sequence elements. Strongly prefers a uridine in the first position of their guide (g1U preference, also named 1U-bias). Not involved in the piRNA amplification loop, also named ping-pong amplification cycle. Acts as an endoribonuclease that cleaves transposon messenger RNAs. Besides their function in transposable elements repression, piRNAs are probably involved in other processes during meiosis such as translation regulation. Probable component of some RISC complex, which mediates RNA cleavage and translational silencing. Also plays a role in the formation of chromatoid bodies and is required for some miRNAs stability. Required to sequester RNF8 in the cytoplasm until late spermatogenesis, RNF8 being released upon ubiquitination and degradation of PIWIL1. {ECO:0000250|UniProtKB:Q9JMB7}, FUNCTION: [Isoform 3]: May be a negative developmental regulator (PubMed:12037681, PubMed:16287078). {ECO:0000269|PubMed:12037681, ECO:0000269|PubMed:16287078}.

Molecular Weight: 98.6 kDa

UniProt: [Q96J94](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: 12 months