

Datasheet for ABIN7555170 RPA1 Protein (AA 1-616) (His tag)



_					
	W	0	rv	10	W

Quantity:	1 mg
Target:	RPA1
Protein Characteristics:	AA 1-616
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This RPA1 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

Purpose:	Custom-made recombinat RPA1 Protein expressed in mammalien cells.
Sequence:	MVGQLSEGAI AAIMQKGDTN IKPILQVINI RPITTGNSPP RYRLLMSDGL NTLSSFMLAT
	QLNPLVEEEQ LSSNCVCQIH RFIVNTLKDG RRVVILMELE VLKSAEAVGV KIGNPVPYNE
	GLGQPQVAPP APAASPAASS RPQPQNGSSG MGSTVSKAYG ASKTFGKAAG PSLSHTSGGT
	QSKVVPIASL TPYQSKWTIC ARVTNKSQIR TWSNSRGEGK LFSLELVDES GEIRATAFNE
	QVDKFFPLIE VNKVYYFSKG TLKIANKQFT AVKNDYEMTF NNETSVMPCE DDHHLPTVQF
	DFTGIDDLEN KSKDSLVDII GICKSYEDAT KITVRSNNRE VAKRNIYLMD TSGKVVTATL
	WGEDADKFDG SRQPVLAIKG ARVSDFGGRS LSVLSSSTII ANPDIPEAYK LRGWFDAEGQ
	ALDGVSISDL KSGGVGGSNT NWKTLYEVKS ENLGQGDKPD YFSSVATVVY LRKENCMYQA
	CPTQDCNKKV IDQQNGLYRC EKCDTEFPNF KYRMILSVNI ADFQENQWVT CFQESAEAIL
	GQNAAYLGEL KDKNEQAFEE VFQNANFRSF IFRVRVKVET YNDESRIKAT VMDVKPVDYR
	EYGRRLVMSI RRSALM Sequence without tag. The proposed Purification-Tag is based on

experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:

Key Benefits:

- Made to order protein from design to production by highly experienced protein experts.
- · Protein expressed in mammalien cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

Grade:

custom-made

Target Details

Target:

RPA1

Alternative Name:

RPA1 (RPA1 Products)

Background:

Replication protein A 70 kDa DNA-binding subunit (RP-A p70) (Replication factor A protein 1) (RF-A protein 1) (Single-stranded DNA-binding protein) [Cleaved into: Replication protein A 70 kDa DNA-binding subunit, N-terminally processed],FUNCTION: As part of the heterotrimeric replication protein A complex (RPA/RP-A), binds and stabilizes single-stranded DNA intermediates, that form during DNA replication or upon DNA stress. It prevents their reannealing and in parallel, recruits and activates different proteins and complexes involved in DNA metabolism (PubMed:27723720, PubMed:27723717). Thereby, it plays an essential role both in DNA replication and the cellular response to DNA damage (PubMed:9430682). In the cellular response to DNA damage, the RPA complex controls DNA repair and DNA damage checkpoint activation. Through recruitment of ATRIP activates the ATR kinase a master regulator of the DNA damage response (PubMed:24332808). It is required for the recruitment

of the DNA double-strand break repair factors RAD51 and RAD52 to chromatin in response to DNA damage (PubMed:17765923). Also recruits to sites of DNA damage proteins like XPA and XPG that are involved in nucleotide excision repair and is required for this mechanism of DNA repair (PubMed:7697716). Also plays a role in base excision repair (BER) probably through interaction with UNG (PubMed:9765279). Also recruits SMARCAL1/HARP, which is involved in replication fork restart, to sites of DNA damage. Plays a role in telomere maintenance (PubMed:17959650, PubMed:34767620). As part of the alternative replication protein A complex, aRPA, binds single-stranded DNA and probably plays a role in DNA repair. Compared to the RPA2-containing, canonical RPA complex, may not support chromosomal DNA replication and cell cycle progression through S-phase. The aRPA may not promote efficient priming by DNA polymerase alpha but could support DNA synthesis by polymerase delta in presence of PCNA and replication factor C (RFC), the dual incision/excision reaction of nucleotide excision repair and RAD51-dependent strand exchange (PubMed:19996105). {ECO:0000269|PubMed:12791985, ECO:0000269|PubMed:17765923, ECO:0000269|PubMed:17959650, ECO:0000269|PubMed:19116208, ECO:0000269|PubMed:19996105, ECO:0000269|PubMed:24332808, ECO:0000269|PubMed:27723717, ECO:0000269|PubMed:27723720, ECO:0000269|PubMed:34767620, ECO:0000269|PubMed:7697716,

Molecular Weight:

68.1 kDa

ECO:0000269|PubMed:9765279}.

UniProt:

P27694

Pathways:

Telomere Maintenance, DNA Damage Repair, Mitotic G1-G1/S Phases, DNA Replication, Chromatin Binding, Synthesis of DNA

ECO:0000269|PubMed:7700386, ECO:0000269|PubMed:9430682,

Application Details

Application Notes:

In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions:

For Research Use only

Handling

Format:

Liquid

Handling

Buffer:	The buffer composition is at the discretion of the manufacturer.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C.	
Expiry Date:	12 months	