

Datasheet for ABIN7555269
POLR2B Protein (AA 1-1174) (His tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	POLR2B
Protein Characteristics:	AA 1-1174
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This POLR2B protein is labelled with His tag.

Product Details

Purpose:	Custom-made recombinant POLR2B Protein expressed in mammalian cells.
Sequence:	MYDADEDMQY DEDDDEITPD LWQEACWIVI SSYFDEKGLV RQQLDSFDEF IQMSVQRIVE DAPPIDLQAE AQHASGEVEE PPRYLLKFEQ IYLSKPTHWE RDGAPSPMMP NEARLRNLTY SAPLYVDITK TVIKEGEEQL QTQHKTFIG KIPIMLRSTY CLLNGLTDRD LCELNECPLD PGGYFIINGS EKVLIQEKM ATNTVYVFAK KDSKYAYTGE CRSCLNSSR PTSTIWVSMML ARGGQGAKKS AIGQRIVATL PYIKQEVPII IVFRALGFVS DRDILEHIY DFEDPEMMEM VKPSLDEAFV IQEQNVALNF IGSRGAKPGV TKEKRIYAK EVLQKEMPLH VGVSDFCETK KAYFLGYMVH RLLLAALGRR ELDDRDHYGN KRLDLAGPLL AFLFRGMFKN LLKEVRIYAQ KFIDRGKDFN LELAIKTRII SDGLKYSLAT GNWGDQKKAH QARAGVSQVL NRLTFASTLS HLRRLNSPIG RDGKLAKPRQ LHNTLWGMVC PAETPEGHAV GLVKNLALMA YISVGSQPSP ILEFLEEWSM ENLEEISPAI IADATKIFVN GCWVGIIHKDP EQLMNTLRKL RRQMDIIVSE VSMIRDIRER EIRIYTDAGR ICRPLLIVEK QKLLLKRHI DQLKEREYNN YSWQDLVASG VVEYIDTLEE ETVMLAMTPD DLQEKEVAYC STYTHCEIHP SMILGVCASI IPFPDHNQSP

Product Details

RNTYQSAMGK QAMGVYITNF HVRMDTLAHV LYYPQKPLVT TRSMEYLRFR ELPAGINSIV
AIASYTGYNQ EDSVIMNRSA VDRGFFRSVF YRSYKEQESK KGFDQEEVFE KPTRETCQGM
RHAIYDKLDD DGLIAPGVRV SGDDVIIGKT VTLPENEDEL ESTNRRYTKR DCSTFLRTSE
TGIVDQVMVT LNQEGYKFKC IRVRSVRIPQ IGDKFASRHG QKGTGCIQYR QEDMPFTCEG
ITPDIINPH AIPSRMTIGH LIECLQGKVS ANKGEIGDAT PFNDAVNVQK ISNLLSDYGY
HLRGNEVLYN GFTGRKITSQ IFIGPTYQR LKHMVDDKIH SRARGPIQIL NRQPMEGRSR
DGGLRFGEME RDCQIAHGAA QFLRERLFEA SDPYQVHVCN LCGIMAIANT RTHTYEGRGC
RNKTQISLVR MPYACKLLFQ ELMSMSIAPR MMSV **Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.**

Specificity: If you are looking for a specific domain and are interested in a partial protein or a different isoform, please contact us regarding an individual offer.

Characteristics: Key Benefits:

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity: > 90 % as determined by Bis-Tris PAGE, anti-tag ELISA, Western Blot and analytical SEC (HPLC)

Grade: custom-made

Target Details

Target: POLR2B

Alternative Name: POLR2B ([POLR2B Products](#))

Target Details

Background:

DNA-directed RNA polymerase II subunit RPB2 (EC 2.7.7.6) (3'-5' exoribonuclease) (EC 3.1.13.-) (DNA-directed RNA polymerase II 140 kDa polypeptide) (DNA-directed RNA polymerase II subunit B) (RNA polymerase II subunit 2) (RNA polymerase II subunit B2) (RNA-directed RNA polymerase II subunit RPB2) (EC 2.7.7.48),FUNCTION: Catalytic core component of RNA polymerase II (Pol II), a DNA-dependent RNA polymerase which synthesizes mRNA precursors and many functional non-coding RNAs using the four ribonucleoside triphosphates as substrates (PubMed:9852112, PubMed:27193682, PubMed:30190596) (By similarity). Pol II-mediated transcription cycle proceeds through transcription initiation, transcription elongation and transcription termination stages. During transcription initiation, Pol II pre-initiation complex (PIC) is recruited to DNA promoters, with focused-type promoters containing either the initiator (Inr) element, or the TATA-box found in cell-type specific genes and dispersed-type promoters that often contain hypomethylated CpG islands usually found in housekeeping genes. Once the polymerase has escaped from the promoter it enters the elongation phase during which RNA is actively polymerized, based on complementarity with the template DNA strand. Transcription termination involves the release of the RNA transcript and polymerase from the DNA (PubMed:9852112, PubMed:27193682, PubMed:30190596). Forms Pol II active center together with the largest subunit POLR2A/RPB1. Appends one nucleotide at a time to the 3' end of the nascent RNA, with POLR2A/RPB1 most likely contributing a Mg(2+)-coordinating DxDGD motif, and POLR2B/RPB2 participating in the coordination of a second Mg(2+) ion and providing lysine residues believed to facilitate Watson-Crick base pairing between the incoming nucleotide and template base. Typically, Mg(2+) ions direct a 5' nucleoside triphosphate to form a phosphodiester bond with the 3' hydroxyl of the preceding nucleotide of the nascent RNA, with the elimination of pyrophosphate. The reversible pyrophosphorolysis can occur at high pyrophosphate concentrations (PubMed:9852112, PubMed:30190596) (By similarity). Can proofread the nascent RNA transcript by means of a 3' -> 5' exonuclease activity. If a ribonucleotide is mis-incorporated, backtracks along the template DNA and cleaves the phosphodiester bond releasing the mis-incorporated 5'-ribonucleotide (PubMed:8381534) (By similarity). {ECO:0000250|UniProtKB:A5PJW8, ECO:0000269|PubMed:27193682, ECO:0000269|PubMed:30190596, ECO:0000269|PubMed:8381534, ECO:0000269|PubMed:9852112}., FUNCTION: RNA-dependent RNA polymerase that catalyzes the extension of a non-coding RNA (ncRNA) at the 3'-end using the four ribonucleoside triphosphates as substrates. An internal ncRNA sequence near the 3'-end serves as a template in a single-round Pol II-mediated RNA polymerization reaction. May decrease the stability of ncRNAs that repress Pol II-mediated gene transcription. {ECO:0000269|PubMed:23395899}.

Target Details

Molecular Weight:	133.9 kDa
UniProt:	P30876
Pathways:	Regulatory RNA Pathways , DNA Damage Repair

Application Details

Application Notes:	We expect the protein to work for functional studies. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months