

## Datasheet for ABIN7555363

# Rubicon Protein (AA 1-972) (His tag)



#### Go to Product page

()	ve	rvi	6	W
$\sim$	v C	1 V I	$\sim$	v v

Quantity:	1 mg
Target:	Rubicon (KIAA0226)
Protein Characteristics:	AA 1-972
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Rubicon protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

Product Details	
Purpose:	Custom-made recombinat RUBCN Protein expressed in mammalien cells.
Sequence:	MRPEGAGMEL GGGEERLPEE SRREHWQLLG NLKTTVEGLV STNSPNVWSK YGGLERLCRD
	MQSILYHGLI RDQACRRQTD YWQFVKDIRW LSPHSALHVE KFISVHENDQ SSADGASERA
	VAELWLQHSL QYHCLSAQLR PLLGDRQYIR KFYTDAAFLL SDAHVTAMLQ CLEAVEQNNP
	RLLAQIDASM FARKHESPLL VTKSQSLTAL PSSTYTPPNS YAQHSYFGSF SSLHQSVPNN
	GSERRSTSFP LSGPPRKPQE SRGHVSPAED QTIQAPPVSV SALARDSPLT PNEMSSSTLT
	SPIEASWVSS QNDSPGDASE GPEYLAIGNL DPRGRTASCQ SHSSNAESSS SNLFSSSSSQ
	KPDSAASSLG DQEGGGESQL SSVLRRSSFS EGQTLTVTSG AKKSHIRSHS DTSIASRGAP
	ESCNDKAKLR GPLPYSGQSS EVSTPSSLYM EYEGGRYLCS GEGMFRRPSE GQSLISYLSE
	QDFGSCADLE KENAHFSISE SLIAAIELMK CNMMSQCLEE EEVEEEDSDR EIQELKQKIR
	LRRQQIRTKN LLPMYQEAEH GSFRVTSSSS QFSSRDSAQL SDSGSADEVD EFEIQDADIR
	RNTASSSKSF VSSQSFSHCF LHSTSAEAVA MGLLKQFEGM QLPAASELEW LVPEHDAPQK

LLPIPDSLPI SPDDGQHADI YKLRIRVRGN LEWAPPRPQI IFNVHPAPTR KIAVAKQNYR
CAGCGIRTDP DYIKRLRYCE YLGKYFCQCC HENAQMAIPS RVLRKWDFSK YYVSNFSKDL
LIKIWNDPLF NVQDINSALY RKVKLLNQVR LLRVQLCHMK NMFKTCRLAK ELLDSFDTVP
GHLTEDLHLY SLNDLTATRK GELGPRLAEL TRAGATHVER CMLCQAKGFI CEFCQNEDDI
IFPFELHKCR TCEECKACYH KACFKSGSCP RCERLQARRE ALARQSLESY LSDYEEEPAE
ALALEAAVLE AT Sequence without tag. The proposed Purification-Tag is based on
experiences with the expression system, a different complexity of the protein could make
another tag necessary. In case you have a special request, please contact us.

#### Characteristics:

Key Benefits:

- Made to order protein from design to production by highly experienced protein experts.
- · Protein expressed in mammalien cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- · State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

## Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

#### Grade:

custom-made

### **Target Details**

Target:	Rubicon (KIAA0226)	
Alternative Name:	RUBCN (KIAA0226 Products)	
Background:	Run domain Beclin-1-interacting and cysteine-rich domain-containing protein (Rubicon) (Beclin-1 associated RUN domain containing protein) (Baron), FUNCTION: Inhibits PIK3C3 activity, under basal conditions negatively regulates PI3K complex II (PI3KC3-C2) function in autophagy.  Negatively regulates endosome maturation and degradative endocytic trafficking and impairs	
	autophagosome maturation process. Can sequester UVRAG from association with a class C	

	Vps complex (possibly the HOPS complex) and negatively regulates Rab7 activation
	(PubMed:20974968, PubMed:21062745). {ECO:0000269 PubMed:20974968,
	ECO:0000269 PubMed:21062745}., FUNCTION: Involved in regulation of pathogen-specific host
	defense of activated macrophages. Following bacterial infection promotes NADH oxidase
	activity by association with CYBA thereby affecting TLR2 signaling and probably other TLR-NOX
	pathways. Stabilizes the CYBA:CYBB NADPH oxidase heterodimer, increases its association
	with TLR2 and its phagosome trafficking to induce antimicrobial burst of ROS and production
	of inflammatory cytokines (PubMed:22423966). Following fungal or viral infection (implicating
	CLEC7A (dectin-1)-mediated myeloid cell activation or RIGI-dependent sensing of RNA viruses)
	negatively regulates pro-inflammatory cytokine production by association with CARD9 and
	sequestering it from signaling complexes (PubMed:22423967).
	{ECO:0000269 PubMed:22423966, ECO:0000269 PubMed:22423967}.
Molecular Weight:	108.6 kDa
UniProt:	Q92622
Pathways:	Autophagy
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a
	guarantee though.
Restrictions:	For Research Use only
	,
Handling	
Handling Format:	Liquid
Format:	Liquid
Format: Buffer:	Liquid  The buffer composition is at the discretion of the manufacturer.
Format:  Buffer:  Handling Advice:	Liquid  The buffer composition is at the discretion of the manufacturer.  Avoid repeated freeze-thaw cycles.