

Datasheet for ABIN755557
SMARCA4 Protein (AA 1-1647) (His tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	SMARCA4
Protein Characteristics:	AA 1-1647
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This SMARCA4 protein is labelled with His tag.

Product Details

Purpose:	Custom-made recombinant SMARCA4 Protein expressed in mammalian cells.
Sequence:	MSTDPPLGG TPRPGSPGP GPSPGAMLGP SPGPSPGSAH SMMGPSPGPP SAGHIPTQG PGGYPQDNMH QMHKPMESMH EKGMSDDPRY NQMKGMGMRS GGHAGMGPPP SPMDQHSQGY PSPLGGSEHA SSPVPASGPS SGPQMSSGPG GAPLDGADPQ ALGQQNRGPT PFNQNLHQL RAQIMAYKML ARGQPLPDHL QMAVQGKRPM PGMQQQMPTL PPPSVSATGP GPGPGPGPGP GPGPAPPNYS RPHGMGGPNM PPPGPGSVPP GMPGQPPGGP PKPWPEGPMA NAAAPTSTPQ KLIPPQTGR PSPAPPAVPP AASPMPPQT QSPGQPAQPA PMVPLHQKQS RITPIQKPRG LDPVEILQER EYRLQARIAH RIQELNLPG SLAGDLRTKA TIELKALRL NFQRQLRQEV VVCMRRDTAL ETALNAKAYK RSKRQSLREA RITEKLEKQQ KIEQERKRRQ KHQEYLSIL QHAKDFKEYH RSVTGKIQKL TKAVATYHAN TEREQKKENE RIEKERMRR MAEDEEGYRK LIDQKKDKRL AYLLQQTDEY VANLTELVRQ HKAAQVAKEK KKKKKKKKAE NAEGQTPAIG PDGEPLDETS QMSDLPVKVI HVESGKILTG TDAPKAGQLE AWLEMNPGYE VAPRSDSEES GSEEEEEEEEE EEQPQAAQPP TLPVEEKKKI PDPDSDDVSE VDAHHIENA

KQDVDDEYGV SQALARGLQS YYAVAHAVTE RVDKQSALMV NGVLKQYQIK GLEWLVSLYN
NNLNGILADE MGLGKTIQTI ALITYLMEHK RINGPFLIIV PLSTLSNWAY EFDKWAPSVV
KVSYKGSPAA RRAFVPQLRS GKFNVLTTY EYIHKHIL AKIRWKYMIV DEGHRMKNHH
CKLTQVLNTH YVAPRRLLT GTPLQNKLPE LWALLNLLP TIFKSCSTFE QWFNAPFAMT
GEKVDLNEEE TILIIRRLHK VLRPFLLRRL KKEVEAQLPE KVEYVIKCDM SALQRVLYRH
MQAKGVLLTD GSEKDKKGGK GTKTLMNTIM QLRKICNHPY MFQHIIESFS EHLGFTGGIV
QGLDLYRASG KFELLDRLP KLRATNHKVL LFCQMTSLMT IMEDYFAYRG FKYLRLDGTT
KAEDRGMLLK TFNEPGSEYF IFLLSTRAGG LGLNLQSADT VIIFDSDWNP HQDLQAQDRA
HRIGQQNEVR VLRLCTVNSV EEKILAAKY KLNVDQKVIQ AGMFDQKSSS HERRAFLQAI
LEHEEQDESR HCSTGSGSAS FAHTAPPPAG VNPDLLEPPL KEEDVPDDE TVNQMIARHE
EEFDLFMRMD LDRRREEARN PKRKPRLMEE DELPSWIKD DAEVERLTCE EEEEKMFGRG
SRHRKEVDYS DSLTEKQWLK AIEEGTLEEI EEEVRQKKSS RKRKRSDAG SSTPTTSTRS
RDKDDESKKQ KKRGRPPAEK LSPNPPNLTK KMKKIVDAVI KYKDSSSGRQ LSEVFIQLPS
RKELPEYYEL IRKPVDFKKI KERIRNHKYR SLNDLEKDVM LLCQNAQTFN LEGSLIYEDS
IVLQSVFTSV RQKIEKEDDS EGESEEEEE GEEGSESES RSVKVKIKLG RKEKAQDRLK
GRRRPSRGS RAKPVVSDDD SSEEQEDRS GSGSEED **Sequence without tag. The proposed
Purification-Tag is based on experiences with the expression system, a different complexity
of the protein could make another tag necessary. In case you have a special request, please
contact us.**

Specificity: If you are looking for a specific domain and are interested in a partial protein or a different isoform, please contact us regarding an individual offer.

Characteristics: Key Benefits:

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein

Product Details

cannot be expressed or purified.

Purity: > 90 % as determined by Bis-Tris PAGE, anti-tag ELISA, Western Blot and analytical SEC (HPLC)

Grade: custom-made

Target Details

Target: SMARCA4

Alternative Name: SMARCA4 ([SMARCA4 Products](#))

Background: Transcription activator BRG1 (EC 3.6.4.-) (ATP-dependent helicase SMARCA4) (BRG1-associated factor 190A) (BAF190A) (Mitotic growth and transcription activator) (Protein BRG-1) (Protein brahma homolog 1) (SNF2-beta) (SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily A member 4),FUNCTION: Involved in transcriptional activation and repression of select genes by chromatin remodeling (alteration of DNA-nucleosome topology). Component of SWI/SNF chromatin remodeling complexes that carry out key enzymatic activities, changing chromatin structure by altering DNA-histone contacts within a nucleosome in an ATP-dependent manner. Component of the CREST-BRG1 complex, a multiprotein complex that regulates promoter activation by orchestrating the calcium-dependent release of a repressor complex and the recruitment of an activator complex. In resting neurons, transcription of the c-FOS promoter is inhibited by SMARCA4-dependent recruitment of a phospho-RB1-HDAC repressor complex. Upon calcium influx, RB1 is dephosphorylated by calcineurin, which leads to release of the repressor complex. At the same time, there is increased recruitment of CREBBP to the promoter by a CREST-dependent mechanism, which leads to transcriptional activation. The CREST-BRG1 complex also binds to the NR2B promoter, and activity-dependent induction of NR2B expression involves the release of HDAC1 and recruitment of CREBBP. Belongs to the neural progenitors-specific chromatin remodeling complex (npBAF complex) and the neuron-specific chromatin remodeling complex (nBAF complex). During neural development, a switch from a stem/progenitor to a postmitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become committed to their adult state. The transition from proliferating neural stem/progenitor cells to postmitotic neurons requires a switch in subunit composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits in neuron-specific complexes (nBAF). The npBAF complex is essential for the self-renewal/proliferative capacity of the multipotent neural stem cells. The nBAF complex along with CREST plays a role

Target Details

regulating the activity of genes essential for dendrite growth. SMARCA4/BAF190A may promote neural stem cell self-renewal/proliferation by enhancing Notch-dependent proliferative signals, while concurrently making the neural stem cell insensitive to SHH-dependent differentiating cues (By similarity). Acts as a corepressor of ZEB1 to regulate E-cadherin transcription and is required for induction of epithelial-mesenchymal transition (EMT) by ZEB1. Binds via DLX1 to enhancers located in the intergenic region between DLX5 and DLX6 and this binding is stabilized by the long non-coding RNA (lncRNA) Evf2 (By similarity). Binds to RNA in a promiscuous manner (By similarity). Binding to RNAs including lncRNA Evf2 leads to inhibition of SMARCA4 ATPase and chromatin remodeling activities (By similarity). In brown adipose tissue, involved in the regulation of thermogenic genes expression (By similarity).
{ECO:0000250|UniProtKB:Q3TKT4, ECO:0000269|PubMed:19571879, ECO:0000269|PubMed:20418909, ECO:0000269|PubMed:29374058, ECO:0000303|PubMed:22952240, ECO:0000303|PubMed:26601204}.

Molecular Weight:	184.6 kDa
UniProt:	P51532
Pathways:	Intracellular Steroid Hormone Receptor Signaling Pathway , Regulation of Intracellular Steroid Hormone Receptor Signaling , Stem Cell Maintenance

Application Details

Application Notes:	We expect the protein to work for functional studies. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months