antibodies .- online.com





anti-CDC7 antibody (AA 151-250) (Cy5.5)



Overview

| Quantity: | 100 μL |
|----------------------|--|
| Target: | CDC7 |
| Binding Specificity: | AA 151-250 |
| Reactivity: | Human, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This CDC7 antibody is conjugated to Cy5.5 |
| Application: | Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human Cdc7 |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human, Rat |
| Predicted Reactivity: | Mouse,Dog,Cow,Pig,Horse |
| Purification: | Purified by Protein A. |

Target Details

| Target: | CDC7 |
|-------------------|----------------------|
| Alternative Name: | Cdc7 (CDC7 Products) |

Target Details

| rarget Details | |
|---------------------|--|
| Background: | Synonyms: Hsk1, CDC7L1, HsCDC7, huCDC7, Cell division cycle 7-related protein kinase, CDC7- |
| | related kinase, CDC7 |
| | Background: Seems to phosphorylate critical substrates that regulate the G1/S phase transition |
| | and/or DNA replication. Can phosphorylates MCM2 and MCM3. |
| Gene ID: | 8317 |
| UniProt: | 000311 |
| Pathways: | Mitotic G1-G1/S Phases, DNA Replication |
| Application Details | |
| Application Notes: | FCM 1:20-100 |
| | IF(IHC-P) 1:50-200 |
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
| | 50 % Glycerol. |
| Preservative: | ProClin |
| | |

Precaution of Use:

Storage Comment:

Storage:

Expiry Date:

handled by trained staff only.

-20 °C

12 months

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.