

Datasheet for ABIN7555814

TOPBP1 Protein (AA 1-1522) (His tag)[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	TOPBP1
Protein Characteristics:	AA 1-1522
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TOPBP1 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Purpose:	Custom-made recombinat TOPBP1 Protein expressed in mammalien cells.
Sequence:	MSRNDKEPFF VKFLKSSDNS KCFFKALES I KEFQSEEYLQ IITEEEALKI KENDRSLYIC DPFSGVVFDH LKKLGCRIVG PQVVIFCMHH QRCVPRAEHP VYNMVMDSVT ISCTSLEKEK REEVHKYVQM MGGRVYRDLN VSVTHLIAGE VGSKKYLVAALNKKPILLPS WIKTLWEKSQ EKKITRYTDI NMEDFKCPIF LGCIICVTGL CGLDRKEVQQ LTVKHGGQYM GQLKMNECTH LIVQEPKGQK YECAKRWNVH CVTTQWFFDS IEKGFCQDES IYKTEPRPEA KTMPNSSTPT SQINTIDSRT LSDVSNISNI NASCVSESI NSLNSKLEPT LENLENLDVS AFQAPEDLLD GCRIYLCGFS GRKLDKLRRL INSGGGVRFN QLNEDVTHVI VGDYDELKQ FWNKSAHRPH VVGAKWLLEC FSKGYMLSEE PYIHANYQPV EIPVSHKPES KAALLKKKNS SFSKKDFAPS EKHEQAEDL LSQYENGSS VVEAKTSEAR PFNDSTHAEP LNDSTHISLQ EENQSSVSHC VPDVSTITEE GLFSQKSFLV LGFSNENESN IANIIKENAG KIMSLLSRTV ADYAVVPLLG CEVEATVGEV VTNTWLVTCTI DYQTLFDPKS NPLFTPVPVM TGMTPLEDCV ISFSQCAGAE

KESLTFLANL LGASVQEYFV RKSNAKKGGMF ASTHLILKER GGSKYEAACK WNLPAVTIAW
LLETARTGKR ADESHFLIEN STKEERSLET EITNGINLNS DTAEHGTR L QTHRKTVVTP
LDMNRFQSKA FRAVVSQHAR QVAASPAVGQ PLQKEPSLHL DTPSKFLSKD KLFKPSFDVK
DALAALETPG RPSQQKRKPS TPLSEVIVKN LQLALANSSR NAVALSASPQ LKEAQSEKKE
APKPLHKVVV CVSKKLSKKQ SELNGIAASL GADYRWSFDE TVTHFIYQGR PNDTNREYKS
VKERGVHIVS EHWLLDCAQE CKHLPESLYP HTYNPKMSLD ISAVQDGRLC NSRLLSAVSS
TKDDEPDPLI LEENDVDNMA TNNKESAPSN GSGKNDKSGV LTQTLEMREN FQKQLQEIMS
ATSIVKPQGQ RTSLSRGCM SASSTPDSTR SARSGRSRVL EALRQSRQTV PDVNTEPSQN
EQIWDPTA REERARLASN LQWPSCPTQY SELQVDIQLN EDSPFQKPLH DSEIAKQAVC
DPGNIRVTEA PKHPISEELE TPIKDSHLIP TPQAPSIAPF LANPPVAPHP REKIITIEET HEELKKQYIF
QLSSLNPQER IDYCHLIEKL GGLVIEKQCF DPTCTHIVVG HPLRNEKYLA SVAAGKWVLH
RSYLEACRTA GHFVQEEDYE WGSSSILDVL TGINVQRRRL ALAAMRWRWK IQRRQESGIV
EGAFSGWKVI LHVDQSREAG FKRLQLQSGGA KVLPGHSVPL FKEATHLFSN LNKLKPDSSG
VNIAEAAAQN VYCLRTEYIA DYLMQESPPH VENYCLPEAI SFIQNNKELG TGLSQKRKAP
TEKNKIKRPR VH **Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.**

Characteristics:

Key Benefits:

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity: > 90 % as determined by Bis-Tris Page, Western Blot

Grade: custom-made

Target Details

Target: TOPBP1

Alternative Name: TOPBP1 ([TOPBP1 Products](#))

Background: DNA topoisomerase 2-binding protein 1 (DNA topoisomerase II-beta-binding protein 1) (TopBP1) (DNA topoisomerase II-binding protein 1),FUNCTION: Scaffold protein that acts as a key protein-protein adapter in DNA replication and DNA repair (PubMed:10498869, PubMed:11395493, PubMed:11714696, PubMed:17575048, PubMed:20545769, PubMed:21777809, PubMed:26811421, PubMed:30898438, PubMed:33592542, PubMed:35597237, PubMed:37674080, PubMed:31135337). Composed of multiple BRCT domains, which specifically recognize and bind phosphorylated proteins, bringing proteins together into functional combinations (PubMed:17575048, PubMed:20545769, PubMed:21777809, PubMed:26811421, PubMed:30898438, PubMed:35597237, PubMed:37674080, PubMed:31135337). Required for DNA replication initiation but not for the formation of pre-replicative complexes or the elongation stages (By similarity). Necessary for the loading of replication factors onto chromatin, including GMNC, CDC45, DNA polymerases and components of the GINS complex (By similarity). Plays a central role in DNA repair by bridging proteins and promoting recruitment of proteins to DNA damage sites (PubMed:30898438, PubMed:35597237, PubMed:37674080). Involved in double-strand break (DSB) repair via homologous recombination in S-phase by promoting the exchange between the DNA replication factor A (RPA) complex and RAD51 (PubMed:26811421, PubMed:35597237). Mechanistically, TOPBP1 is recruited to DNA damage sites in S-phase via interaction with phosphorylated HTATSF1, and promotes the loading of RAD51, thereby facilitating RAD51 nucleofilaments formation and RPA displacement, followed by homologous recombination (PubMed:35597237). Involved in microhomology-mediated end-joining (MMEJ) DNA repair by promoting recruitment of polymerase theta (POLQ) to DNA damage sites during mitosis (PubMed:37674080). MMEJ is an alternative non-homologous end-joining (NHEJ) machinery that takes place during mitosis to repair DSBs in DNA that originate in S-phase (PubMed:37674080). Recognizes and binds POLQ phosphorylated by PLK1, enabling its recruitment to DSBs for subsequent repair (PubMed:37674080). Involved in G1 DNA damage checkpoint by acting as a molecular adapter that couples TP53BP1 and the 9-1-1 complex (PubMed:31135337). In response to DNA damage, triggers the recruitment of checkpoint signaling proteins on chromatin, which activate the CHEK1 signaling pathway and block S-phase progression (PubMed:16530042, PubMed:21777809). Acts as an activator of the kinase activity of ATR (PubMed:16530042, PubMed:21777809). Also required for chromosomal stability when DSBs occur during mitosis by forming filamentous assemblies that bridge MDC1 and tether broken chromosomes during mitosis (PubMed:30898438). Together with CIP2A,

Target Details

plays an essential role in the response to genome instability generated by the presence of acentric chromosome fragments derived from shattered chromosomes within micronuclei (PubMed:35121901, PubMed:35842428, PubMed:37165191, PubMed:37316668). Micronuclei, which are frequently found in cancer cells, consist of chromatin surrounded by their own nuclear membrane: following breakdown of the micronuclear envelope, a process associated with chromothripsis, the CIP2A-TOPBP1 complex tethers chromosome fragments during mitosis to ensure clustered segregation of the fragments to a single daughter cell nucleus, facilitating re-ligation with limited chromosome scattering and loss (PubMed:37165191, PubMed:37316668). Recruits the SWI/SNF chromatin remodeling complex to E2F1-responsive promoters, thereby down-regulating E2F1 activity and inhibiting E2F1-dependent apoptosis during G1/S transition and after DNA damage (PubMed:12697828, PubMed:15075294). {ECO:0000250|UniProtKB:Q800K6, ECO:0000269|PubMed:10498869, ECO:0000269|PubMed:11395493, ECO:0000269|PubMed:11714696, ECO:0000269|PubMed:12697828, ECO:0000269|PubMed:15075294, ECO:0000269|PubMed:16530042, ECO:0000269|PubMed:17575048, ECO:0000269|PubMed:20545769, ECO:0000269|PubMed:21777809, ECO:0000269|PubMed:26811421, ECO:0000269|PubMed:30898438, ECO:0000269|PubMed:31135337, ECO:0000269|PubMed:33592542, ECO:0000269|PubMed:35121901, ECO:0000269|PubMed:35597237, ECO:0000269|PubMed:35842428, ECO:0000269|PubMed:37165191, ECO:0000269|PubMed:37316668, ECO:0000269|PubMed:37674080}.

Molecular Weight: 170.7 kDa

UniProt: [Q92547](#)

Pathways: [Chromatin Binding](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions: For Research Use only

Handling

Format: Liquid

Handling

Buffer: The buffer composition is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: 12 months