

Datasheet for ABIN7555885
TRPV4 Protein (AA 1-871) (His tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	TRPV4
Protein Characteristics:	AA 1-871
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TRPV4 protein is labelled with His tag.

Product Details

Purpose:	Custom-made recombinant TRPV4 Protein expressed in mammalian cells.
Sequence:	MADSSEGPRA GPGEVAELPG DESGTPGGEA FPLSSLANLF EGEDGSLSPS PADASRPAGP GDGRPNLRMK FQGAFRKGVP NPIDLLESTL YESSVVPGPK KAPMDSLFDY GTYRHHSSDN KRWRKKIEK QPQSPKAPAP QPPPILKVFN RPILFDIVSR GSTADLDGLL PFLLTTHKKRL TDEEFREPST GKTCLPKALL NLSNGRNDTI PVLLDIAERT GNMREFINSP FRDIYYRGQT ALHIAIERRC KHYVELLVAQ GADVHAQARG RFFQPKDEGG YFYFGELPLS LAACTNQPHI VNYLTENPHK KADMRRQDSR GNTVLHALVA IADNTRENTK FVTKMYDLLL LKCARLFPDS NLEAVLNNDG LSPLMMAAKT GKIGIFQHII RREVTDEDTR HLSRKFKDWA YGPVYSSLYD LSSLDTCGEE ASVLEILVYN SKIENRHEML AVEPINELLR DKWRKFGAVS FYINVVSYLC AMVIFTLTAY YQPLEGTPPY PYRTTVDYLR LAGEVITLFT GVLFFFTNIK DLFMKKCPGV NSLFDIGSFQ LLYFIYSVLV IVSAALYLAG IEAYLAVMVF ALVLGWMNAL YFTRGLKLTG TYSIMIQKIL FKDLFRLLV YLLFMIGYAS ALVSLLNPCA NMKVCNEDQT NCTVPTYPSC RDSETFSTFL LDLFKLTIGM GDLEMLSSTK YPVVFIILLV TYIILTFVLL LNMLIALMGE

Product Details

TVGQVSKESK HIWKLQWATT ILDIERSFPV FLRKAFRSGE MVTVGKSSDG TPDRRWCFRV
DEVNWSHWNQ NLGIINEDPG KNETYQYYGF SHTVGRLRRD RWSSVVPVV ELNKNSNPDE
VVVPLDSMGN PRCDGHQQGY PRKWRTDDAP L **Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.**

Specificity: If you are looking for a specific domain and are interested in a partial protein or a different isoform, please contact us regarding an individual offer.

Characteristics: Key Benefits:

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity: > 90 % as determined by Bis-Tris PAGE, anti-tag ELISA, Western Blot and analytical SEC (HPLC)

Grade: custom-made

Target Details

Target: TRPV4

Alternative Name: TRPV4 ([TRPV4 Products](#))

Background: Transient receptor potential cation channel subfamily V member 4 (TrpV4) (Osm-9-like TRP channel 4) (OTRPC4) (Transient receptor potential protein 12) (TRP12) (Vanilloid receptor-like channel 2) (Vanilloid receptor-like protein 2) (VRL-2) (Vanilloid receptor-related osmotically-activated channel) (VR-OAC),FUNCTION: Non-selective calcium permeant cation channel involved in osmotic sensitivity and mechanosensitivity (PubMed:16293632, PubMed:18826956,

PubMed:18695040, PubMed:29899501, PubMed:22526352, PubMed:23136043). Activation by exposure to hypotonicity within the physiological range exhibits an outward rectification (PubMed:18826956, PubMed:18695040, PubMed:29899501). Also activated by heat, low pH, citrate and phorbol esters (PubMed:16293632, PubMed:18826956, PubMed:18695040, PubMed:25256292, PubMed:20037586, PubMed:21964574). Increase of intracellular Ca(2+) potentiates currents. Channel activity seems to be regulated by a calmodulin-dependent mechanism with a negative feedback mechanism (PubMed:12724311, PubMed:18826956). Promotes cell-cell junction formation in skin keratinocytes and plays an important role in the formation and/or maintenance of functional intercellular barriers (By similarity). Acts as a regulator of intracellular Ca(2+) in synoviocytes (PubMed:19759329). Plays an obligatory role as a molecular component in the nonselective cation channel activation induced by 4-alpha-phorbol 12,13-didecanoate and hypotonic stimulation in synoviocytes and also regulates production of IL-8 (PubMed:19759329). Together with PKD2, forms mechano- and thermosensitive channels in cilium (PubMed:18695040). Negatively regulates expression of PPARGC1A, UCP1, oxidative metabolism and respiration in adipocytes (By similarity). Regulates expression of chemokines and cytokines related to pro-inflammatory pathway in adipocytes (By similarity). Together with AQP5, controls regulatory volume decrease in salivary epithelial cells (By similarity). Required for normal development and maintenance of bone and cartilage (PubMed:26249260). In its inactive state, may sequester DDX3X at the plasma membrane. When activated, the interaction between both proteins is affected and DDX3X relocalizes to the nucleus (PubMed:29899501). In neurons of the central nervous system, could play a role in triggering voluntary water intake in response to increased sodium concentration in body fluid (By similarity). {ECO:0000250|UniProtKB:Q9EPK8, ECO:0000269|PubMed:11025659, ECO:0000269|PubMed:12724311, ECO:0000269|PubMed:16293632, ECO:0000269|PubMed:18587396, ECO:0000269|PubMed:18695040, ECO:0000269|PubMed:18826956, ECO:0000269|PubMed:19759329, ECO:0000269|PubMed:20037586, ECO:0000269|PubMed:21964574, ECO:0000269|PubMed:23136043, ECO:0000269|PubMed:25256292, ECO:0000269|PubMed:26249260, ECO:0000269|PubMed:29899501}., FUNCTION: [Isoform 1]: Non-selective calcium permeant cation channel involved in osmotic sensitivity and mechanosensitivity. Activation by exposure to hypotonicity within the physiological range exhibits an outward rectification. Also activated by phorbol esters. Has the same channel activity as isoform 1, and is activated by the same stimuli. {ECO:0000269|PubMed:16293632}., FUNCTION: [Isoform 5]: Non-selective calcium permeant cation channel involved in osmotic sensitivity and mechanosensitivity. Activation by exposure to hypotonicity within the physiological range exhibits an outward rectification. Also activated by phorbol esters. Has the

Target Details

same channel activity as isoform 1, and is activated by the same stimuli.

{ECO:0000269|PubMed:16293632}, FUNCTION: [Isoform 2]: Lacks channel activity, due to impaired oligomerization and intracellular retention. {ECO:0000269|PubMed:16293632}, FUNCTION: [Isoform 4]: Lacks channel activity, due to impaired oligomerization and intracellular retention. {ECO:0000269|PubMed:16293632}, FUNCTION: [Isoform 6]: Lacks channel activity, due to impaired oligomerization and intracellular retention. {ECO:0000269|PubMed:16293632}, FUNCTION: (Microbial infection) Facilitates hepatitis C virus (HCV) replication, possibly through its action on DDX3X. {ECO:0000269|PubMed:29899501}, FUNCTION: (Microbial infection) Facilitates Dengue virus (DENV) replication, possibly through its action on DDX3X. {ECO:0000269|PubMed:29899501}, FUNCTION: (Microbial infection) Facilitates Zika virus (ZIKV) replication, possibly through its action on DDX3X. {ECO:0000269|PubMed:29899501}.

Molecular Weight: 98.3 kDa

UniProt: [Q9HBA0](#)

Pathways: [Hormone Transport, Cell-Cell Junction Organization](#)

Application Details

Application Notes: We expect the protein to work for functional studies. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: 12 months