

Datasheet for ABIN7556862 NME2 Protein (AA 1-152) (His tag)



Overview

Quantity:	1 mg
Target:	NME2
Protein Characteristics:	AA 1-152
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This NME2 protein is labelled with His tag.

Custom-made recombinant Nme2 Protein expressed in mammalian cells.
MANLERTFIA IKPDGVQRGL VGEIIKRFEQ KGFRLVAMKF LRASEEHLKQ HYIDLKDRPF
FPGLVKYMNS GPVVAMVWEG LNVVKTGRVM LGETNPADSK PGTIRGDFCI QVGRNIIHGS
DSVESAEKEI HLWFKPEELI DYKSCAHDWV YE Sequence without tag. The proposed
Purification-Tag is based on experiences with the expression system, a different complexity
of the protein could make another tag necessary. In case you have a special request, please
contact us.
If you are looking for a specific domain and are interested in a partial protein or a different
isoform, please contact us regarding an individual offer.
Key Benefits:
Made to order protein - from design to production - by highly experienced protein experts.
Protein expressed in mammalian cells and purified in one-step affinity chromatography
The optimized expression system ensures reliability for intracellular, secreted and

transmembrane proteins.

• State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris PAGE, anti-tag ELISA, Western Blot and analytical SEC (HPLC)

Grade:

custom-made

Target Details

Target:	NME2
Alternative Name:	Nme2 (NME2 Products)
Background:	Nucleoside diphosphate kinase B (NDK B) (NDP kinase B) (EC 2.7.4.6) (Histidine protein kinase
	NDKB) (EC 2.7.13.3) (P18) (nm23-M2),FUNCTION: Major role in the synthesis of nucleoside
	triphosphates other than ATP. The ATP gamma phosphate is transferred to the NDP beta
	phosphate via a ping-pong mechanism, using a phosphorylated active-site intermediate (By
	similarity). Negatively regulates Rho activity by interacting with AKAP13/LBC. Acts as a
	transcriptional activator of the MYC gene, binds DNA non-specifically. Binds to both single-
	stranded guanine- and cytosine-rich strands within the nuclease hypersensitive element (NHE)
	III(1) region of the MYC gene promoter. Does not bind to duplex NHE III(1). Has G-quadruplex
	(G4) DNA-binding activity, which is independent of its nucleotide-binding and kinase activity.
	Binds both folded and unfolded G4 with similar low nanomolar affinities. Stabilizes folded G4s
	regardless of whether they are prefolded or not. Exhibits histidine protein kinase activity (By
	similarity). {ECO:0000250 UniProtKB:P22392, ECO:0000250 UniProtKB:P36010,
	ECO:0000269 PubMed:20884616}.
Molecular Weight:	17.4 kDa
UniProt:	Q01768

Application Details

Application Notes:	We expect the protein to work for functional studies. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months