

Datasheet for ABIN7557132 SDHAF1 Protein (AA 1-118) (His tag)



Go to Product page

\sim					
	1//	Р	rv	I P	۱۸/

Quantity:	1 mg
Target:	SDHAF1
Protein Characteristics:	AA 1-118
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This SDHAF1 protein is labelled with His tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB)
Product Details	
Purpose:	Custom-made recombinat Sdhaf1 Protein expressed in mammalien cells.
Sequence:	MSRPSRLQRQ VLSLYRELLR AGRGTPGAEA RVRAEFRQHA SLPRTDVLRI EYLYRRGRRQ
	LQLLRSGHAT AMGTFVRPRG PAEEPGDATA PGTRLDDGGA PKNSCEDTGA RETRSDGR
	Sequence without tag. The proposed Purification-Tag is based on experiences with the
	expression system, a different complexity of the protein could make another tag necessary.
	In case you have a special request, please contact us.
Characteristics:	Key Benefits:
	Made to order protein - from design to production - by highly experienced protein experts.
	Protein expressed in mammalien cells and purified in one-step affinity chromatography
	 The optimized expression system ensures reliability for intracellular, secreted and
	transmembrana prataina
	transmembrane proteins.

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

Grade:

custom-made

guarantee though.

Target Details

Target:	SDHAF1	
Alternative Name:	Sdhaf1 (SDHAF1 Products)	
Background:	Succinate dehydrogenase assembly factor 1, mitochondrial (SDH assembly factor 1) (SDHAF1) (LYR motif-containing protein 8),FUNCTION: Plays an essential role in the assembly of succinate dehydrogenase (SDH), an enzyme complex (also referred to as respiratory complex II) that is a component of both the tricarboxylic acid (TCA) cycle and the mitochondrial electron transport chain, and which couples the oxidation of succinate to fumarate with the reduction of ubiquinone (coenzyme Q) to ubiquinol. Promotes maturation of the iron-sulfur protein subunit Sdhb of the SDH catalytic dimer, protecting it from the deleterious effects of oxidants. May act together with SDHAF3. Contributes to iron-sulfur cluster incorporation into SDHB by binding to SDHB and recruiting the iron-sulfur transfer complex formed by HSC20, HSPA9 and ISCU through direct binding to HSC20. {ECO:0000250 UniProtKB:A6NFY7}.	
Molecular Weight:	13.1 kDa	
UniProt:	Q3U276	
Application Details		
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a	

Application Details

Expiry Date:

Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.

12 months