

Datasheet for ABIN7557232

CDK10 Protein (AA 1-360) (His tag)



\sim				
()\	/e	r\/		٨
() 1	v C.	ı vı	\Box	ΙV

DK10
1-360
ouse
EK-293 Cells
combinant
is CDK10 protein is labelled with His tag.
estern Blotting (WB), SDS-PAGE (SDS)
ol Ek

Product Details

Purpose:	Custom-made recombinat Cdk10 Protein expressed in mammalien cells.
Sequence:	MAEVDLESDQ IRLKCIRKEG FFTVPPEHRL GRCRSVKEFE KLNRIGEGTY GIVYRARDTQ
	TDEIVALKKV RMDKEKDGIP ISSLREITLL LRLRHPNIVE LKEVVVGNHL ESIFLVMGYC
	EQDLASLLEN MPTPFSEAQV KCIMLQVLRG LQYLHRNFII HRDLKVSNLL MTDKGCVKTA
	DFGLARAYGV PVKPMTPKVV TLWYRAPELL LGTTTQTTSI DMWAVGCILA ELLAHKPLLP
	GTSEIHQIDL IVQLLGTPSE NIWPGFSKLP LAGQYSLRKQ PYNNLKHKFP WLSEAGLRLL
	NFLFMYDPKK RATSGDCLES SYFKEKPLPC EPELMPTFPH HRNKRAAPAA AEGQSKRCRP
	Sequence without tag. The proposed Purification-Tag is based on experiences with the
	expression system, a different complexity of the protein could make another tag necessary.
	In case you have a special request, please contact us.
Characteristics:	Key Benefits:

- · Made to order protein from design to production by highly experienced protein experts.
- · Protein expressed in mammalien cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

Grade:

custom-made

Target Details

Target:	CDK10
Alternative Name:	Cdk10 (CDK10 Products)
Background:	Cyclin-dependent kinase 10 (EC 2.7.11.22) (Cell division protein kinase 10),FUNCTION: Cyclin-dependent kinase that phosphorylates the transcription factor ETS2 (in vitro) and positively controls its proteasomal degradation (in cells). Involved in the regulation of actin cytoskeleton organization through the phosphorylation of actin dynamics regulators such as PKN2. Is a negative regulator of ciliogenesis through phosphorylation of PKN2 and promotion of RhoA signaling. {ECO:0000250 UniProtKB:Q15131}.
Molecular Weight:	41.0 kDa
UniProt:	Q3UMM4

Application Details

Application Notes:

In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Application Details

Storage Comment:

Expiry Date:

Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C

Store at -80°C.

12 months