

Datasheet for ABIN7557618
NRF2 Protein (AA 1-597) (His tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	NRF2 (NFE2L2)
Protein Characteristics:	AA 1-597
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This NRF2 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Purpose:	Custom-made recombinat Nfe2l2 Protein expressed in mammalian cells.
Sequence:	MMDLELPPPG LQSQQDMDLI DILWRQDIDL GVSREVFDFS QRQKDYELEK QKKLEKERQE QLQKEQEKAFAQFQLDEET GEFLPIQPAQ HIQTDTSRSA SYSQVAHIPK QDALYFEDCM QLLAETFPFV DDHESLALDI PSHAESSVFT APHQAQSLNS SLEAAMTDLS SIEQDMEQVW QELFSIPELQ CLNTENKQLA DTTAVPSPEA TLTEMDSNYH FYSSISSLEK EVGNCGPHFL HGFEDSFSSI LSTDDASQLT SLDSNPTLNT DFGDEFYSAF IAEPDGGGSM PSSAAISQSL SELLDGTIEG CDLSLCKAFN PKHAEGTMEF NDSDSGISLN TSPSRASPEH SVESSYGDGP PPGFSSEME ELDSAPGSVK QNGPKAQAHA SPGDTVQPLS PAQGHSAAPMR ESQCENNTKK EVPVSPGHQK APFTKDKHSS RLEAHLTRDE LRAKALHIPF PVEKIINLPV DDFNEMMSKE QFNEAQLALI RDIRRRGKNK VAAQNCRKRK LENIVELEQD LGHLKDEREK LLREKGENDR NLHLLKRRLS TLYLEVFSML RDEDGKPYSP SEYSLQQTRD GNVFLVPKSK KPDTKKK Sequence without tag. The proposed Purification-Tag is based on experiences with the expression

system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:	<p>Key Benefits:</p> <ul style="list-style-type: none">• Made to order protein - from design to production - by highly experienced protein experts.• Protein expressed in mammalian cells and purified in one-step affinity chromatography• The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.• State-of-the-art algorithm used for plasmid design (Gene synthesis). <p>This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.</p> <p>If you are not interested in a full length protein, please contact us for individual protein fragments.</p> <p>The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.</p>
------------------	---

Purity:	> 90 % as determined by Bis-Tris Page, Western Blot
Grade:	custom-made

Target Details

Target:	NRF2 (NFE2L2)
Alternative Name:	Nfe2l2 (NFE2L2 Products)
Background:	<p>Nuclear factor erythroid 2-related factor 2 (NF-E2-related factor 2) (NFE2-related factor 2) (Nuclear factor, erythroid derived 2, like 2),FUNCTION: Transcription factor that plays a key role in the response to oxidative stress: binds to antioxidant response (ARE) elements present in the promoter region of many cytoprotective genes, such as phase 2 detoxifying enzymes, and promotes their expression, thereby neutralizing reactive electrophiles (PubMed:9240432, PubMed:9887101, PubMed:12032331, PubMed:14517554, PubMed:31398338). In normal conditions, ubiquitinated and degraded in the cytoplasm by the BCR(KEAP1) complex (PubMed:15282312, PubMed:15367669, PubMed:15581590). In response to oxidative stress, electrophile metabolites inhibit activity of the BCR(KEAP1) complex, promoting nuclear accumulation of NFE2L2/NRF2, heterodimerization with one of the small Maf proteins and binding to ARE elements of cytoprotective target genes (PubMed:12032331). The</p>

Target Details

NFE2L2/NRF2 pathway is also activated in response to selective autophagy: autophagy promotes interaction between KEAP1 and SQSTM1/p62 and subsequent inactivation of the BCR(KEAP1) complex, leading to NFE2L2/NRF2 nuclear accumulation and expression of cytoprotective genes (PubMed:20421418, PubMed:20173742). May also be involved in the transcriptional activation of genes of the beta-globin cluster by mediating enhancer activity of hypersensitive site 2 of the beta-globin locus control region (By similarity). Also plays an important role in the regulation of the innate immune response. It is a critical regulator of the innate immune response and survival during sepsis by maintaining redox homeostasis and restraint of the dysregulation of pro-inflammatory signaling pathways like MyD88-dependent and -independent and TNF-alpha signaling (PubMed:16585964). Suppresses macrophage inflammatory response by blocking pro-inflammatory cytokine transcription and the induction of IL6 (PubMed:27211851). Binds to the proximity of pro-inflammatory genes in macrophages and inhibits RNA Pol II recruitment. The inhibition is independent of the Nrf2-binding motif and reactive oxygen species level (PubMed:27211851). Represses antiviral cytosolic DNA sensing by suppressing the expression of the adapter protein STING1 and decreasing responsiveness to STING1 agonists while increasing susceptibility to infection with DNA viruses (By similarity). {ECO:0000250|UniProtKB:Q16236, ECO:0000269|PubMed:12032331, ECO:0000269|PubMed:14517554, ECO:0000269|PubMed:15282312, ECO:0000269|PubMed:15367669, ECO:0000269|PubMed:15581590, ECO:0000269|PubMed:16585964, ECO:0000269|PubMed:20173742, ECO:0000269|PubMed:20421418, ECO:0000269|PubMed:27211851, ECO:0000269|PubMed:31398338, ECO:0000269|PubMed:9240432, ECO:0000269|PubMed:9887101}.

Molecular Weight: 66.9 kDa

UniProt: [Q60795](#)

Pathways: [ER-Nucleus Signaling](#), [Negative Regulation of intrinsic apoptotic Signaling](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months