

## Datasheet for ABIN7557645

## CYP7B1 Protein (AA 1-507) (His tag)



Go to Product page

()	ve	r\/i	Δ	۱۸/
$\circ$	V C	1 V		v v

Quantity:	1 mg
Target:	CYP7B1
Protein Characteristics:	AA 1-507
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CYP7B1 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

Product Details		
Purpose:	Custom-made recombinat Cyp7b1 Protein expressed in mammalien cells.	
Sequence:	MQGATTLDAA SPGPLALLGL LFAATLLLSA LFLLTRRTRR PREPPLIKGW LPYLGMALKF	
	FKDPLTFLKT LQRQHGDTFT VFLVGKYITF VLNPFQYQYV TKNPKQLSFQ KFSSRLSAKA	
	FSVKKLLTDD DLNEDVHRAY LLLQGKPLDA LLETMIQEVK ELFESQLLKI TDWNTERIFA	
	FCGSLVFEIT FATLYGKILA GNKKQIISEL RDDFFKFDDM FPYLVSDIPI QLLRNEESMQ	
	KKIIKCLTSE KVAQMQGQSK IVQERQDLLK RYYRHDDPEI GAHHLGFLWA SLANTIPAMF	
	WAMYYILRHP EAMEALRDEI DSFLQSTGQK KGPGISVHFT REQLDSLVCL ESTILEVLRL	
	CSYSSIIREV QEDMNLSLES KSFSLRKGDF VALFPPLIHN DPEIFDAPKE FRFDRFIEDG	
	KKKSTFFKGG KKLKTYVMPF GLGTSKCPGR YFAVNEMKLL LIMLLTYFDL EIIDRKPIGL	
	NHSRMFLGIQ HPDSAVSFRY KAKSWRS Sequence without tag. The proposed Purification-	
	Tag is based on experiences with the expression system, a different complexity of the	
	protein could make another tag necessary. In case you have a special request, please	

## contact us. Characteristics: Key Benefits: Made to order protein - from design to production - by highly experienced protein experts. Protein expressed in mammalien cells and purified in one-step affinity chromatography · The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins. • State-of-the-art algorithm used for plasmid design (Gene synthesis). This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein. If you are not interested in a full length protein, please contact us for individual protein fragments. The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified. > 90 % as determined by Bis-Tris Page, Western Blot Purity: Grade: custom-made **Target Details** CYP7B1 Target: Alternative Name: Cyp7b1 (CYP7B1 Products) Background: Cytochrome P450 7B1 (24-hydroxycholesterol 7-alpha-hydroxylase) (EC 1.14.14.26) (25/26hydroxycholesterol 7-alpha-hydroxylase) (EC 1.14.14.29) (3-hydroxysteroid 7-alpha hydroxylase) (Hippocampal transcript 1 protein) (HCT-1) (Oxysterol 7-alphahydroxylase), FUNCTION: A cytochrome P450 monooxygenase involved in the metabolism of endogenous oxysterols and steroid hormones, including neurosteroids (PubMed:9295351, PubMed:9144166, PubMed:10748047, PubMed:11067870). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (CPR, NADPH-ferrihemoprotein reductase). Catalyzes the hydroxylation of carbon hydrogen bonds of steroids with a preference for 7-alpha position. Usually metabolizes steroids carrying a hydroxy

group at position 3, functioning as a 3-hydroxy steroid 7-alpha hydroxylase (PubMed:9295351,

PubMed:9144166, PubMed:10748047, PubMed:11067870). Hydroxylates oxysterols, including

Target Details		
	25-hydroxycholesterol and (25R)-cholest-5-ene-3beta,26-diol toward 7-alpha hydroxy	
	derivatives, which may be transported to the liver and converted to bile acids	
	(PubMed:9295351, PubMed:9144166, PubMed:10748047, PubMed:11067870). Via its product	
	7-alpha,25-dihydroxycholesterol, a ligand for the chemotactic G protein-coupled receptor	
	GPR183/EBI2, regulates B cell migration in germinal centers of lymphoid organs, thus guiding	
	efficient maturation of plasma B cells and overall antigen-specific humoral immune response	
	(PubMed:22999953). 7-alpha hydroxylates neurosteroids, including 3beta-hydroxyandrost-5-en-	
	17-one (dehydroepiandrosterone) and pregnenolone, both involved in hippocampus-associated	
	memory and learning (PubMed:9144166). Metabolizes androstanoids toward 6- or 7-alpha	
	hydroxy derivatives (By similarity). {ECO:0000250 UniProtKB:075881,	
	ECO:0000269 PubMed:10748047, ECO:0000269 PubMed:11067870,	
	ECO:0000269 PubMed:22999953, ECO:0000269 PubMed:9144166,	
	ECO:0000269 PubMed:9295351}.	
Molecular Weight:	58.5 kDa	
UniProt:	Q60991	
Pathways:	Intracellular Steroid Hormone Receptor Signaling Pathway, Steroid Hormone Biosynthesis,	
	Regulation of Intracellular Steroid Hormone Receptor Signaling	
Application Details		
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies	

Restrictions:

Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months

guarantee though.

For Research Use only

as well. As the protein has not been tested for functional studies yet we cannot offer a