

Datasheet for ABIN7557792 NDUFA4 Protein (AA 1-82) (Fc Tag)



Overview

Quantity:	1 mg
Target:	NDUFA4
Protein Characteristics:	AA 1-82
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This NDUFA4 protein is labelled with Fc Tag.

Product Details

Custom-made recombinant Ndufa4 Protein expressed in mammalian cells.
MLRQILGQAK KHPSLIPLFV FIGAGGTGAA LYVMRLALFN PDVSWDRKNN PEPWNKLGPN
EQYKFYSVNV DYSKLKKEGP DF Sequence without tag. The proposed Purification-Tag is
based on experiences with the expression system, a different complexity of the protein
could make another tag necessary. In case you have a special request, please contact us.
If you are looking for a specific domain and are interested in a partial protein or a different
isoform, please contact us regarding an individual offer.
Key Benefits:
Made to order protein - from design to production - by highly experienced protein experts.
 Protein expressed in mammalian cells and purified in one-step affinity chromatography
 The optimized expression system ensures reliability for intracellular, secreted and
transmembrane proteins.
 State-of-the-art algorithm used for plasmid design (Gene synthesis).
-

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris PAGE, anti-tag ELISA, Western Blot and analytical SEC (HPLC)

Grade:

custom-made

Target Details

Target:	NDUFA4
Alternative Name:	Ndufa4 (NDUFA4 Products)
Background:	Cytochrome c oxidase subunit NDUFA4,FUNCTION: Component of the cytochrome c oxidase,
	the last enzyme in the mitochondrial electron transport chain which drives oxidative
	phosphorylation. The respiratory chain contains 3 multisubunit complexes succinate
	dehydrogenase (complex II, CII), ubiquinol-cytochrome c oxidoreductase (cytochrome b-c1
	complex, complex III, CIII) and cytochrome c oxidase (complex IV, CIV), that cooperate to
	transfer electrons derived from NADH and succinate to molecular oxygen, creating an
	electrochemical gradient over the inner membrane that drives transmembrane transport and
	the ATP synthase. Cytochrome c oxidase is the component of the respiratory chain that
	catalyzes the reduction of oxygen to water. Electrons originating from reduced cytochrome c in
	the intermembrane space (IMS) are transferred via the dinuclear copper A center (CU(A)) of
	subunit 2 and heme A of subunit 1 to the active site in subunit 1, a binuclear center (BNC)
	formed by heme A3 and copper B (CU(B)). The BNC reduces molecular oxygen to 2 water
	molecules unsing 4 electrons from cytochrome c in the IMS and 4 protons from the
	mitochondrial matrix. NDUFA4 is required for complex IV maintenance.
	{ECO:0000250 UniProtKB:000483}.
Molecular Weight:	9.3 kDa
UniProt:	Q62425

Application Details

Application Notes:	We expect the protein to work for functional studies. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months