

## Datasheet for ABIN7557982

# CCDC111 Protein (AA 1-537) (His tag)



## Overview

Quantity:	1 mg
Target:	CCDC111
Protein Characteristics:	AA 1-537
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CCDC111 protein is labelled with His tag.

Product Details	
Purpose:	Custom-made recombinant Primpol Protein expressed in mammalian cells.
Sequence:	MLRKWEARVK QIEERASHYE RKPLSSVYRP RLAKPEEPSS IWKLFHRQNQ AFNFVKSCKE
	SVHVFALECK RGNGQRIYLV TSYAQLWFYY KTRKTLLHCY EVIPENAVCK LYFDLEFNKL
	ANPGADGKMM VALLIQHVCK ALEEFYNVQC SAEDVFNLDS STEEKFSRHL IFQLHNVAFK
	DNRHAGNFVR KILQPALHLI AEDDEAKVPE AVGQDASGFS VTPLKQEISE AREKVGLPKQ
	CDPDLSFLVV KNHMGEKCLF VDLGVYTKNR NFRLYQSSKI GKCVSLEVAE DNRFIPKQSK
	DISEENQYFL SSLVSNVRFS DTLRVLTCHP SQTKRKRAEC FNSTGTSVES IEGFQGSPYP
	EVDQFVLSLV NKHDIKGGIR RWNYFFPEEL LVYDICKYRW CENIGRAHKS NNIMILVDLK
	NEVWYQKCHD PVCKAQNFKS TCSPLPTEVS LLFLLKDEDF TSGETDDTST SLTKDSQTPP
	SCNLSAGGLS AAAWDDEDDA LFLEATEDAE FADAADKSLG SMDDIPDELI IEALQNS <b>Sequence</b>
	without tag. The proposed Purification-Tag is based on experiences with the expression
	system, a different complexity of the protein could make another tag necessary. In case you
	have a special request, please contact us.

#### **Product Details**

Product Details	
Specificity:	If you are looking for a specific domain and are interested in a partial protein or a different
	isoform, please contact us regarding an individual offer.
Characteristics:	Key Benefits:
	Made to order protein - from design to production - by highly experienced protein experts.
	Protein expressed in mammalian cells and purified in one-step affinity chromatography
	The optimized expression system ensures reliability for intracellular, secreted and
	transmembrane proteins.
	State-of-the-art algorithm used for plasmid design (Gene synthesis).
	This protein is a made-to-order protein and will be made for the first time for your order. Our
	experts in the lab try to ensure that you receive soluble protein.
	If you are not interested in a full length protein, please contact us for individual protein
	fragments.
	The big advantage of ordering our made-to-order proteins in comparison to ordering custom
	made proteins from other companies is that there is no financial obligation in case the protein
	cannot be expressed or purified.
Purity:	> 90 % as determined by Bis-Tris PAGE, anti-tag ELISA, Western Blot and analytical SEC (HPLC
Grade:	custom-made
Target Details	
Target:	CCDC111
Alternative Name:	Primpol (CCDC111 Products)
Background:	DNA-directed primase/polymerase protein (EC 2.7.7),FUNCTION: DNA primase and DNA
	polymerase required to tolerate replication-stalling lesions by bypassing them
	(PubMed:26926109, PubMed:29073063). Required to facilitate mitochondrial and nuclear
	replication fork progression by initiating de novo DNA synthesis using dNTPs and acting as a
	error-prone DNA polymerase able to bypass certain DNA lesions (By similarity). Shows a high
	capacity to tolerate DNA damage lesions such as 80x0G and abasic sites in DNA (By similarit
	dapatity to tolerate brist damage lesions such as boxod and abasic sites in bina (by similarity

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn | International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com | Page 2/3 | Product datasheet for ABIN7557982 | 03/28/2025 | Copyright antibodies-online. All rights reserved.

Provides different translesion synthesis alternatives when DNA replication is stalled: able to

quadruplexes, to allow DNA replication to continue (By similarity). Can also realign primers

ahead of 'unreadable lesions' such as abasic sites and 6-4 photoproduct (6-4 pyrimidine-

pyrimidinone), thereby skipping the lesion (By similarity). Also able to incorporate nucleotides

synthesize DNA primers downstream of lesions, such as ultraviolet (UV) lesions, R-loops and G-

opposite DNA lesions such as 80xoG, like a regular translesion synthesis DNA polymerase (By similarity). Also required for reinitiating stalled forks after UV damage during nuclear DNA replication (By similarity). Required for mitochondrial DNA (mtDNA) synthesis and replication, by reinitiating synthesis after UV damage or in the presence of chain-terminating nucleotides (PubMed:29073063). Prevents APOBEC family-mediated DNA mutagenesis by repriming downstream of abasic site to prohibit error-prone translesion synthesis (PubMed:26926109). Has non-overlapping function with POLH (By similarity). In addition to its role in DNA damage response, also required to maintain efficient nuclear and mitochondrial DNA replication in unperturbed cells (By similarity). {ECO:0000250|UniProtKB:Q96LW4, ECO:0000269|PubMed:26926109, ECO:0000269|PubMed:29073063}.

Molecular Weight:

61.3 kDa

UniProt:

**Q6P1E7** 

### **Application Details**

**Application Notes:** 

We expect the protein to work for functional studies. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions:

For Research Use only

## Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months