antibodies .- online.com





anti-FE65 antibody (pSer347)



0 -	4.4	Prod	400		
1-()	10	Pron	ויאוו	nac	\Box
\circ	LO.	1 100	uot	Duc	

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	FE65 (APBB1)
Binding Specificity:	pSer347
Reactivity:	Mouse, Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FE65 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human APBB1 around the phosphorylation site of Ser347 [SL(p-S)PE]
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.
Target Details	
Target:	FE65 (APBB1)

Target Details

Alternative Name:	APBB1 (APBB1 Products)		
Background:	Synonyms: Adaptor protein FE65a2, Amyloid beta A4 precursor protein binding family B		
	member 1, Amyloid beta A4 precursor protein binding family B, Amyloid beta A4 precursor		
	protein binding family B member 1, Amyloid beta precursor protein binding family B member 1		
	APBB 1, APBB1, FE 65, Fe65 protein, MGC 9072, MGC9072, Protein Fe65, RIR, Stat like protein,		
	Fe-65 Protein, APBB1_HUMAN.		
	Background: The protein encoded by this gene is a member of the Fe65 protein family. It is an		
	adaptor protein localized in the nucleus. It interacts with the Alzheimer's disease amyloid		
	precursor protein (APP), transcription factor CP2/LSF/LBP1 and the low-density lipoprotein		
	receptor-related protein. APP functions as a cytosolic anchoring site that can prevent the gene		
	product's nuclear translocation. This encoded protein could play an important role in the		
	pathogenesis of Alzheimer's disease. It is thought to regulate transcription. Also it is observed		
	to block cell cycle progression by downregulating thymidylate synthase expression. Multiple		
	alternatively spliced transcript variants encoding different isoforms have been described for		
	this gene.		
Gene ID:	322		
Pathways:	Positive Regulation of Response to DNA Damage Stimulus		
Application Details			
Application Notes:	WB 1:300-5000		
	ELISA 1:500-1000		
	IHC-P 1:200-400		
	IHC-F 1:100-500		
	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.		

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months