

Datasheet for ABIN7559917
HDAC8 Protein (AA 1-377) (His tag)



[Go to Product page](#)

Overview

| | |
|-------------------------------|--|
| Quantity: | 1 mg |
| Target: | HDAC8 |
| Protein Characteristics: | AA 1-377 |
| Origin: | Mouse |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This HDAC8 protein is labelled with His tag. |
| Application: | SDS-PAGE (SDS), Western Blotting (WB) |

Product Details

| | |
|------------------|---|
| Purpose: | Custom-made recombinat Hdac8 Protein expressed in mammalien cells. |
| Sequence: | MEMPEEPANS GHSLPPVYIY SPEYVSICDS LVKVPKRASM VHSLIEAYAL HKQMRIVKPK VASMEEMATF HTDAYLQHLQ KVSQEGDEDH PDSIEYGLGY DCPATEGIFD YAAAIGGGTI TAAQCLIDGK CKVAINWSGG WHHAKKDEAS GFCYLNDAVL GILRLRRKFD RILYVDLDLH HGDGVEDAFS FTSKVMTVSL HKFSPGFFPG TGDMSDVGLG KGRYYSVNVP IQDGIQDEKY YHICESVLKE VYQAFNPKAV VLQLGADTIA GDPMCSFNMT PVGIGKCLKY VLQWQLATLI LGGGGYNLAN TARCWTYLTG VILGKTLSS EIPDHEFFTAY GPDYVLEITP SCRPRDRNEPH RIQQILNYIK GNLKHVV Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us. |
| Characteristics: | Key Benefits: |

Product Details

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity: > 90 % as determined by Bis-Tris Page, Western Blot

Grade: custom-made

Target Details

Target: HDAC8

Alternative Name: Hdac8 ([HDAC8 Products](#))

Background: Histone deacetylase 8 (HD8) (EC 3.5.1.98) (Protein deacetylase HDAC8) (EC 3.5.1.-) (Protein deacetylase HDAC8) (EC 3.5.1.-),FUNCTION: Histone deacetylase that catalyzes the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Also involved in the deacetylation of cohesin complex protein SMC3 regulating release of cohesin complexes from chromatin. May play a role in smooth muscle cell contractility. In addition to protein deacetylase activity, also has protein-lysine deacylase activity: acts as a protein deacetylase by mediating deacetylation ((2E)-butenoyl) of histones. {ECO:0000250|UniProtKB:Q9BY41}.

Molecular Weight: 41.8 kDa

UniProt: [Q8VH37](#)

Pathways: [Cellular Glucan Metabolic Process](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: 12 months
