

Datasheet for ABIN7560473
COX6C Protein (AA 1-76) (Fc Tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	COX6C
Protein Characteristics:	AA 1-76
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This COX6C protein is labelled with Fc Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Purpose:	Custom-made recombinat Cox6c Protein expressed in mammalien cells.
Sequence:	MSSGALLPKP QMRGLLAKRL RVHIAGAFIV ALGVAAAYKF GVAEPRKKAY AEFYRNYDSM KDFEEMRKAG IFQSAK Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.
Characteristics:	<p>Key Benefits:</p> <ul style="list-style-type: none"> • Made to order protein - from design to production - by highly experienced protein experts. • Protein expressed in mammalien cells and purified in one-step affinity chromatography • The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins. • State-of-the-art algorithm used for plasmid design (Gene synthesis). <p>This protein is a made-to-order protein and will be made for the first time for your order. Our</p>

experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:	> 90 % as determined by Bis-Tris Page, Western Blot
---------	---

Grade:	custom-made
--------	-------------

Target Details

Target:	COX6C
---------	-------

Alternative Name:	Cox6c (COX6C Products)
-------------------	--

Background:	<p>Cytochrome c oxidase subunit 6C (Cytochrome c oxidase polypeptide VIc),FUNCTION: Component of the cytochrome c oxidase, the last enzyme in the mitochondrial electron transport chain which drives oxidative phosphorylation. The respiratory chain contains 3 multisubunit complexes succinate dehydrogenase (complex II, CII), ubiquinol-cytochrome c oxidoreductase (cytochrome b-c1 complex, complex III, CIII) and cytochrome c oxidase (complex IV, CIV), that cooperate to transfer electrons derived from NADH and succinate to molecular oxygen, creating an electrochemical gradient over the inner membrane that drives transmembrane transport and the ATP synthase. Cytochrome c oxidase is the component of the respiratory chain that catalyzes the reduction of oxygen to water. Electrons originating from reduced cytochrome c in the intermembrane space (IMS) are transferred via the dinuclear copper A center (CU(A)) of subunit 2 and heme A of subunit 1 to the active site in subunit 1, a binuclear center (BNC) formed by heme A3 and copper B (CU(B)). The BNC reduces molecular oxygen to 2 water molecules using 4 electrons from cytochrome c in the IMS and 4 protons from the mitochondrial matrix. {ECO:0000250 UniProtKB:P04038}.</p>
-------------	---

Molecular Weight:	8.5 kDa
-------------------	---------

UniProt:	Q9CPQ1
----------	------------------------

Pathways:	Proton Transport
-----------	----------------------------------

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months