

Datasheet for ABIN7560616
PNRC2 Protein (AA 1-140) (His tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	PNRC2
Protein Characteristics:	AA 1-140
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This PNRC2 protein is labelled with His tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB)

Product Details

Purpose:	Custom-made recombinat Pnrc2 Protein expressed in mammalien cells.
Sequence:	MGGGERYNIP DPQSRNASKN QEQQNRQKSK DQNSSQTKIA HKKKERGHGY NPAAAAWQAM QNGGKTKSLS NNSNWNAGLS SPSLLFKSQA SQNYAGAKFS EPPSPSVLPK PPSHWVHVSL NPSDKETMTF QLKTLLKVQV Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:	<p>Key Benefits:</p> <ul style="list-style-type: none">• Made to order protein - from design to production - by highly experienced protein experts.• Protein expressed in mammalien cells and purified in one-step affinity chromatography• The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.• State-of-the-art algorithm used for plasmid design (Gene synthesis).
------------------	--

Product Details

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity: > 90 % as determined by Bis-Tris Page, Western Blot

Grade: custom-made

Target Details

Target: PNRC2

Alternative Name: Pnrc2 ([PNRC2 Products](#))

Background: Proline-rich nuclear receptor coactivator 2,FUNCTION: Involved in nonsense-mediated mRNA decay (NMD) by acting as a bridge between the mRNA decapping complex and the NMD machinery (By similarity). May act by targeting the NMD machinery to the P-body and recruiting the decapping machinery to aberrant mRNAs (By similarity). Required for UPF1/RENT1 localization to the P-body (By similarity). Plays a role in glucocorticoid receptor-mediated mRNA degradation by interacting with the glucocorticoid receptor NR3C1 in a ligand-dependent manner when it is bound to the 5' UTR of target mRNAs and recruiting the RNA helicase UPF1 and the mRNA-decapping enzyme DCP1A, leading to RNA decay (By similarity). Also acts as a nuclear receptor coactivator. May play a role in controlling the energy balance between energy storage and energy expenditure (PubMed:17971453). {ECO:0000250|UniProtKB:Q9NPJ4, ECO:0000269|PubMed:17971453}.

Molecular Weight: 15.4 kDa

UniProt: [Q9CR73](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Application Details

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: 12 months