

# Datasheet for ABIN7561171

## TBC1D20 Protein (AA 1-402) (His tag)



## Overview

| Quantity:                     | 1 mg   |
|-------------------------------|--|
| Target:                       | TBC1D20  |
| Protein Characteristics:      | AA 1-402                                       |
| Origin:                       | Mouse  |
| Source:                       | HEK-293 Cells                                  |
| Protein Type:                 | Recombinant                                    |
| Purification tag / Conjugate: | This TBC1D20 protein is labelled with His tag. |

| Product Details |   |
|-----------------|---|
| Purpose:        | Custom-made recombinant Tbc1d20 Protein expressed in mammalian cells.                           |
| Sequence:       | MALRPSKGDG SAGRWDRGAG KADFNAKRKK KVAEIHQALN SDPIDLAALR RMAISEGGLL                               |
|                 | TDEIRCQVWP KLLNVNTSEP PPVSRKDLRD MSKDYQQVLL DVRRSLRRFP PGMPDEQREG                               |
|                 | LQEELIDIIL LVLDRNPQLH YYQGYHDIVV TFLLVVGERL ATSLVEKLST HHLRDFMDPT                               |
|                 | MDNTKHILNY LMPIIDQVSP ELHDFMQSAE VGTIFALSWL ITWFGHVLMD FRHVVRLYDF                               |
|                 | FLACHPLMPI YFAAVIVLYR EQEVLDCDCD MASVHHLLSQ IPQDLPYETL ISRAGDLFVQ                               |
|                 | FPPSELAREA AAQQEAERTA ASTFKDFELA STQQRPDMVL RQRFRGLLRP EARTKDVLTK                               |
|                 | PRTNRFVKLA VMGLTVALGA AALAVVKSAL EWAPKFQLQL FP Sequence without tag. The                        |
|                 | proposed Purification-Tag is based on experiences with the expression system, a different       |
|                 | complexity of the protein could make another tag necessary. In case you have a special          |
|                 | request, please contact us.   |
| Specificity:    | If you are looking for a specific domain and are interested in a partial protein or a different |
|                 | isoform, please contact us regarding an individual offer.                                       |

### **Product Details**

| Characteristics:  | Key Benefits:  |
|---|--|
|   | <ul> <li>Made to order protein - from design to production - by highly experienced protein experts.</li> <li>Protein expressed in mammalian cells and purified in one-step affinity chromatography</li> <li>The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.</li> <li>State-of-the-art algorithm used for plasmid design (Gene synthesis).</li> </ul> |
|   | This protein is a made-to-order protein and will be made for the first time for your order. Our  |
|   | experts in the lab try to ensure that you receive soluble protein.   |
|   | If you are not interested in a full length protein, please contact us for individual protein fragments.  |
|   | The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.   |
| Purity:   | > 90 % as determined by Bis-Tris PAGE, anti-tag ELISA, Western Blot and analytical SEC (HPLC)  |
| Grade:  | custom-made  |
| Target Details  |  |
| Target:   | TBC1D20  |
|   |  |
| Alternative Name:   | Tbc1d20 (TBC1D20 Products)   |
| Alternative Name:  Background:                                | Tbc1d20 (TBC1D20 Products)  TBC1 domain family member 20,FUNCTION: GTPase-activating protein specific for Rab1 and Rab2 small GTPase families for which it can accelerate the intrinsic GTP hydrolysis rate by more than five orders of magnitude (By similarity). Involved in maintaining endoplasmic reticulum structure (By similarity). {ECO:0000250 UniProtKB:Q96BZ9}.                                      |
|   | TBC1 domain family member 20,FUNCTION: GTPase-activating protein specific for Rab1 and Rab2 small GTPase families for which it can accelerate the intrinsic GTP hydrolysis rate by more than five orders of magnitude (By similarity). Involved in maintaining endoplasmic   |
| Background:   | TBC1 domain family member 20,FUNCTION: GTPase-activating protein specific for Rab1 and Rab2 small GTPase families for which it can accelerate the intrinsic GTP hydrolysis rate by more than five orders of magnitude (By similarity). Involved in maintaining endoplasmic reticulum structure (By similarity). {ECO:0000250 UniProtKB:Q96BZ9}.  |
| Background:  Molecular Weight:                                | TBC1 domain family member 20,FUNCTION: GTPase-activating protein specific for Rab1 and Rab2 small GTPase families for which it can accelerate the intrinsic GTP hydrolysis rate by more than five orders of magnitude (By similarity). Involved in maintaining endoplasmic reticulum structure (By similarity). {ECO:0000250 UniProtKB:Q96BZ9}.  45.8 kDa  |
| Background:  Molecular Weight:  UniProt:                      | TBC1 domain family member 20,FUNCTION: GTPase-activating protein specific for Rab1 and Rab2 small GTPase families for which it can accelerate the intrinsic GTP hydrolysis rate by more than five orders of magnitude (By similarity). Involved in maintaining endoplasmic reticulum structure (By similarity). {ECO:0000250 UniProtKB:Q96BZ9}.  45.8 kDa  |
| Background:  Molecular Weight:  UniProt:  Application Details | TBC1 domain family member 20,FUNCTION: GTPase-activating protein specific for Rab1 and Rab2 small GTPase families for which it can accelerate the intrinsic GTP hydrolysis rate by more than five orders of magnitude (By similarity). Involved in maintaining endoplasmic reticulum structure (By similarity). {ECO:0000250 UniProtKB:Q96BZ9}.  45.8 kDa  Q9D9I4  |

## Handling

| Format:          | Liquid   |
|------------------|--|
| Buffer:          | The buffer composition is at the discretion of the manufacturer. |
| Handling Advice: | Avoid repeated freeze-thaw cycles.                               |
| Storage:         | -80 °C   |
| Storage Comment: | Store at -80°C.  |
| Expiry Date:     | 12 months  |