

Datasheet for ABIN7561858

NLR Family, Pyrin Domain Containing 1B (NLRP1B) (AA 1-1177) protein (His tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	NLR Family, Pyrin Domain Containing 1B (NLRP1B)
Protein Characteristics:	AA 1-1177
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	His tag
Application:	SDS-PAGE (SDS), Western Blotting (WB)

Product Details

Purpose:	Custom-made recombinat Nlrp1b Protein expressed in mammalian cells.
Sequence:	MEESQYKQEH NKKVAQDEGQ EDKDTIFETI EAIEAKLMEL KTNPESTFNY GIFPEVYMNQ GEEILYPAWS LKEENLFQTF KSLRFLQKLC PRGSGNLVKK SWYPCVPEEG GHIINIQLDF GPNIGTQKEP QLVIIEGAAG IGKSTLARQV KRAWMEGELY RDHFQHVFFF SCRELAQCKK LSLAELITQG QDVPTAPINQ ILSHPEKLLF ILDGIDPAW VLADQNPELC LYWSQTQPVH TLLGSLGKS ILPEASFLLT TRTTALQKFI PSLPQSCQVE VLGFSDFEQE IYIKYFAKQ IFGIKALMMV ESNPVLLTLC EVPWVCWLVC NCLKKQMEQG GDVSLTSQTT TAICLKYISL TIPVHHMRTQ LRALCSLAAE GIWKRRTLFS ESDLCKQGLD EDAVAIFLKT GVLQKQASSL SYFAHLCLQ EFFASMSCIL EDSEERHGDM EMDRIVETLV ERYGRQNLFE APTVRFLFGL LSKEGLKEME KLFSCSLPGK TKLKLLWHIL GKSQPHQPPC LGLLHCLYEN QDMKLLTHVM HDLQGTIVPD TDDITHTVLQ TNVKHLVVRT DMELMVVTFQ IQFCSHMRSQQLNMEGQQGY ALTAPRMVLY RWTPITNASW KILFYNLKFN SNLEGLDLSG NPLSYSQVAVQY LCDAMIYPGC

Product Details

QLKTLWLVEC GLTPTYCSLL ASVLSACSSL RELDLQLNDL CDDGVRMLCE GLRNRACNLR
ILRLDLYSLS AQVITELRTL EENNLKLHIS SIWMPQMMVP TENMDEEDIL TSFKQQRQQS
GANPMEILGT EEDFWGPIGP VATEVVYRER NLYRVQLPMA GSYHCPSTRL HFVVTRAVTI
EIEFCAWSQF LDKTPLQQSH MIVGPLFDIK AEQGAVTAVY LPHFVSLKDT KASTFDFKVA
HFQEHEGMVLE TPDRVKPGYT VLKNPSFSPM GVVLRIPAA RHFIPITSIT LIYYRVNQEE
VTLHLYLVPN DCTIQKAIDD EEMKFQFVRI NKPPPVDNLF IGSRYIVSGS ENLEITPKEL
ELCYRSSKEF QLFSEIYVGN MGSEIKLQIK NKKHMKLIWE ALLKPGDLRP ALPRIAQALK
DAPSLHFMD QHREQLVARV TSDVPLLDKL HGLVLNEESY EAVRAENTNQ DKMRKLFNLS
RSWSRACKDL FYQALKETHP HLVMDLLEKS GGVSLGS **Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.**

Characteristics:

Key Benefits:

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

Grade:

custom-made

Target Details

Target:

NLR Family, Pyrin Domain Containing 1B (NLRP1B)

Alternative Name:

Nlrp1b ([NLRP1B Products](#))

Background:

NACHT, LRR and PYD domains-containing protein 1b allele 2 (EC 3.4.-.-) [Cleaved into: NACHT,

LRR and PYD domains-containing protein 1b, C-terminus (Nlrp1b1-CT), NACHT, LRR and PYD domains-containing protein 1b, N-terminus (Nlrp1b1-NT)],FUNCTION: Acts as the sensor component of the Nlrp1b inflammasome, which mediates inflammasome activation in response to various pathogen-associated signals, leading to subsequent pyroptosis (By similarity). Inflammasomes are supramolecular complexes that assemble in the cytosol in response to pathogens and other damage-associated signals and play critical roles in innate immunity and inflammation (By similarity). Acts as a recognition receptor (PRR): recognizes specific pathogens and other damage-associated signals: in response to pathogen-associated signals, the N-terminal part of Nlrp1b is degraded by the proteasome, releasing the cleaved C-terminal part of the protein (NACHT, LRR and PYD domains-containing protein 1b, C-terminus), which polymerizes to initiate the formation of the inflammasome complex: the inflammasome directly recruits pro-caspase-1 (proCASP1) independently of PYCARD/ASC and promotes caspase-1 (CASP1) activation, which subsequently cleaves and activates inflammatory cytokines IL1B and IL18 and gasdermin-D (GSDMD), leading to pyroptosis (By similarity). In the absence of GSDMD expression, the Nlrp1b inflammasome is able to recruit and activate CASP8, leading to activation of gasdermin-E (GSDME) (By similarity). Activation of Nlrp1b inflammasome is also required for HMGB1 secretion, the active cytokines and HMGB1 stimulate inflammatory responses (By similarity). Contrary to Nlrp1b allele 1, allele 2 is not activated by Bacillus anthracis lethal toxin (PubMed:16429160, PubMed:21170303, PubMed:24492532). {ECO:0000250|UniProtKB:Q2LKW6, ECO:0000269|PubMed:16429160, ECO:0000269|PubMed:21170303, ECO:0000269|PubMed:24492532}., FUNCTION: [NACHT, LRR and PYD domains-containing protein 1b allele 2]: Constitutes the precursor of the Nlrp1b inflammasome, which mediates autoproteolytic processing within the FIIND domain to generate the N-terminal and C-terminal parts, which are associated non-covalently in absence of pathogens and other damage-associated signals. {ECO:0000250|UniProtKB:Q2LKW6}., FUNCTION: [NACHT, LRR and PYD domains-containing protein 1b, N-terminus]: Regulatory part that prevents formation of the Nlrp1b inflammasome: in absence of pathogens and other damage-associated signals, interacts with the C-terminal part of Nlrp1b (NACHT, LRR and PYD domains-containing protein 1b, C-terminus), preventing activation of the Nlrp1b inflammasome. In response to pathogen-associated signals, this part is ubiquitinated by the N-end rule pathway and degraded by the proteasome, releasing the cleaved C-terminal part of the protein, which polymerizes and forms the Nlrp1b inflammasome. {ECO:0000250|UniProtKB:Q2LKW6}., FUNCTION: [NACHT, LRR and PYD domains-containing protein 1b, C-terminus]: Constitutes the active part of the Nlrp1b inflammasome. In absence of pathogens and other damage-associated signals, interacts with the N-terminal part of Nlrp1b (NACHT, LRR and PYD domains-containing protein 1b, N-terminus), preventing activation of the Nlrp1b inflammasome.

Target Details

In response to pathogen-associated signals, the N-terminal part of Nlrp1b is degraded by the proteasome, releasing this form, which polymerizes to form the Nlrp1b inflammasome complex: the Nlrp1b inflammasome complex then directly recruits pro-caspase-1 (proCASP1) and promotes caspase-1 (CASP1) activation, leading to gasdermin-D (GSDMD) cleavage and subsequent pyroptosis. {ECO:0000250|UniProtKB:Q2LKW6}.

Molecular Weight: 134.2 kDa

UniProt: [A1Z198](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: 12 months