

Datasheet for ABIN7562230 SLC22A1 Protein (AA 1-556) (His tag)



Go to Product page

\sim			
()\	/ e	rVI	iew

Quantity:	1 mg
Target:	SLC22A1
Protein Characteristics:	AA 1-556
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This SLC22A1 protein is labelled with His tag.

Product Details

Purpose:	Custom-made recombinant Slc22a1 Protein expressed in mammalian cells.
Sequence:	MPTVDDVLEH VGEFGWFQKQ AFLLLCLISA SLAPIYVGIV FLGFTPDHHC RSPGVAELSQ
	RCGWSPAEEL NYTVPGLGSA GEASFLSQCM KYEVDWNQST LDCVDPLSSL AANRSHLPLS
	PCEHGWVYDT PGSSIVTEFN LVCGDAWKVD LFQSCVNLGF FLGSLVVGYI ADRFGRKLCL
	LVTTLVTSLS GVLTAVAPDY TSMLLFRLLQ GMVSKGSWVS GYTLITEFVG SGYRRTTAIL
	YQVAFTVGLV GLAGVAYAIP DWRWLQLAVS LPTFLFLLYY WFVPESPRWL LSQKRTTQAV
	RIMEQIAQKN RKVPPADLKM MCLEEDASER RSPSFADLFR TPSLRKHTLI LMYLWFSCAV
	LYQGLIMHVG ATGANLYLDF FYSSLVEFPA AFIILVTIDR IGRIYPIAAS NLVAGAACLL
	MIFIPHELHW LNVTLACLGR MGATIVLQMV CLVNAELYPT FIRNLGMMVC SALCDLGGIF
	TPFMVFRLME VWQALPLILF GVLGLSAGAV TLLLPETKGV ALPETIEEAE NLGRRKSKAK
	ENTIYLQVQT GKSPHT Sequence without tag. The proposed Purification-Tag is based on
	experiences with the expression system, a different complexity of the protein could make
	another tag necessary. In case you have a special request, please contact us.

Product Details

Product Details	
Specificity:	If you are looking for a specific domain and are interested in a partial protein or a different isoform, please contact us regarding an individual offer.
Characteristics:	Key Benefits:
Grial acteristics.	 Made to order protein - from design to production - by highly experienced protein experts. Protein expressed in mammalian cells and purified in one-step affinity chromatography The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins. State-of-the-art algorithm used for plasmid design (Gene synthesis).
	This protein is a made-to-order protein and will be made for the first time for your order. Our
	experts in the lab try to ensure that you receive soluble protein.
	If you are not interested in a full length protein, please contact us for individual protein fragments.
	The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.
Purity:	> 90 % as determined by Bis-Tris PAGE, anti-tag ELISA, Western Blot and analytical SEC (HPLC)
Grade:	custom-made
Target Details	
Target:	SLC22A1
Alternative Name:	Slc22a1 (SLC22A1 Products)
Background:	Solute carrier family 22 member 1 (Organic cation transporter 1) (mOCT1),FUNCTION: Electrogenic voltage-dependent transporter that mediates the transport of a variety of organic cations such as endogenous bioactive amines, cationic drugs and xenobiotics (PubMed:10216142, PubMed:12176030, PubMed:11463829, PubMed:23458604, PubMed:24961373). Functions as a pH - and Na(+)-independent, bidirectional transporter (By similarity). Cation cellular uptake or release is driven by the electrochemical potential (i.e.
	oartty). Outlon ochular aptaric or release to arriver by the electrochemical potential (i.e.

membrane potential and concentration gradient) and substrate selectivity (By similarity).

Hydrophobicity is a major requirement for recognition in polyvalent substrates and inhibitors (PubMed:23458604). Primarily expressed in the basolateral membrane of hepatocytes and

proximal tubules and involved in the uptake and disposition of cationic compounds from the

blood by hepatic and renal clearance (By similarity). Most likely functions as an uptake carrier in

enterocytes contributing to the intestinal elimination of organic cations from the systemic circulation (PubMed:11463829, PubMed:24961373). Transports endogenous monoamines such as N-1-methylnicotinamide (NMN), guanidine, neurotransmitters dopamine, serotonin, noradrenaline, adrenaline and histamine, and quaternary ammonium compound such as choline (PubMed:24961373, PubMed:35469921). Also transports natural polyamines such as spermidine, agmatine and putrescine at low affinity, but relatively high turnover (PubMed:23458604). Involved in the hepatic and intestinal uptake of the vitamin B1/thiamine, hence regulating hepatic lipid and energy metabolism (PubMed:24961373). Contributes to the influx and efflux of fatty acid carriers carnitines and acylcarnitines across the basolateral membrane of hepatocytes, from the liver to the systemic circulation and inversely and may be involved in regulating the systemic availability of hepatic acylcarnitines (PubMed:28942964, PubMed:34040533). Also capable of transporting non-amine endogenous compounds such as prostaglandin E2 (PGE2) and prostaglandin F2-alpha (PGF2-alpha) (By similarity). May contribute to the transport of cationic compounds in testes across the blood-testis-barrier (By similarity). Also mediates the uptake of xenobiotics tributylmethylammonium (TBuMA), quinidine, N-methyl-quinine (NMQ), N-methyl-quinidine (NMQD) N-(4,4-azo-n-pentyl)quinuclidine (APQ), azidoprocainamide methoiodide (AMP), N-(4,4-azo-n-pentyl)-21deoxyajmalinium (APDA) and 4-(4-(dimethylamino)styryl)-N-methylpyridinium (ASP) (PubMed:11463829). {ECO:0000250|UniProtKB:015245, ECO:0000250|UniProtKB:Q63089, ECO:0000269|PubMed:10216142, ECO:0000269|PubMed:11463829, ECO:0000269|PubMed:12176030, ECO:0000269|PubMed:23458604, ECO:0000269|PubMed:24961373, ECO:0000269|PubMed:28942964, ECO:0000269|PubMed:34040533, ECO:0000269|PubMed:35469921}.

Molecular Weight:

61.5 kDa

UniProt:

008966

Pathways:

Hormone Transport

Application Details

Application Notes:

We expect the protein to work for functional studies. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions:

For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months