

Datasheet for ABIN7562248
HDAC1 Protein (AA 1-482) (His tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	HDAC1
Protein Characteristics:	AA 1-482
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This HDAC1 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Purpose:	Custom-made recombinat Hdac1 Protein expressed in mammalien cells.
Sequence:	<p>MAQTQGTKRK VCYYYDGDVG NYYYGQGHM KPHRIRMTHN LLLNYGLYRK MEIYRPHKAN AEEMTKYHSD DYIKFLRSIR PDNMSEYSKQ MQRFNVEDC PVFDGLFEFC QLSTGGSVAS AVKLNKQQTD IAVNWAGGLH HAKKSEASGF CYVNDIVLAI LELLKYHQRV LYIDIDIHHG DGVVEAFYTT DRVMTVSFHK YGEYFPGTGD LRDIGAGKGK YYAVNYPLRD GIDDESYEAI FKPVMSKVME MFQPSAVVLQ CGSDSLSGDR LGCFNLTIKG HAKCVEFVKS FNLPLMLLGG GGYTIRNVAR CWTYETAVAL DTEIPNELPY NDYFEYFGPD FKLHISPSNM TNQNTNEYLE KIKQRLFENL RMLPHAPGVQ MQAIPEDAIP EESGDEDEED PDKRISICSS DKRIACEEEF SDSDEEGEGG RKNSSNFKKA KRVKTEDEKE KDPEEKKEVT EEEKTKKEKP EAKGVKEEVK LA</p> <p>Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.</p>

Product Details

Characteristics:

Key Benefits:

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

Grade:

custom-made

Target Details

Target:

HDAC1

Alternative Name:

Hdac1 ([HDAC1 Products](#))

Background:

Histone deacetylase 1 (HD1) (EC 3.5.1.98) (Protein deacetylase HDAC1) (EC 3.5.1.-) (Protein deacetylase HDAC1) (EC 3.5.1.-),FUNCTION: Histone deacetylase that catalyzes the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4) (PubMed:10615135, PubMed:15542849, PubMed:21960634, PubMed:30279482). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events (PubMed:10615135, PubMed:15542849, PubMed:21960634). Histone deacetylases act via the formation of large multiprotein complexes (PubMed:10615135, PubMed:21960634). Acts as a component of the histone deacetylase NuRD complex which participates in the remodeling of chromatin (By similarity). As part of the SIN3B complex is recruited downstream of the constitutively active genes transcriptional start sites through interaction with histones and mitigates histone acetylation and RNA polymerase II progression within transcribed regions contributing to the regulation of transcription (By similarity). Also functions as a deacetylase for non-histone

Target Details

targets, such as NR1D2, RELA, SP1, SP3, STAT3 and TSHZ3 (By similarity). Deacetylates SP proteins, SP1 and SP3, and regulates their function (By similarity). Component of the BRG1-RB1-HDAC1 complex, which negatively regulates the CREST-mediated transcription in resting neurons (By similarity). Upon calcium stimulation, HDAC1 is released from the complex and CREBBP is recruited, which facilitates transcriptional activation (By similarity). Deacetylates TSHZ3 and regulates its transcriptional repressor activity (By similarity). Deacetylates 'Lys-310' in RELA and thereby inhibits the transcriptional activity of NF-kappa-B (By similarity).

Deacetylates NR1D2 and abrogates the effect of KAT5-mediated relieving of NR1D2 transcription repression activity (By similarity). Component of a RCOR/GFI/KDM1A/HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development (PubMed:17707228). Involved in CIART-mediated transcriptional repression of the circadian transcriptional activator: CLOCK-BMAL1 heterodimer (PubMed:15226430, PubMed:24736997). Required for the transcriptional repression of circadian target genes, such as PER1, mediated by the large PER complex or CRY1 through histone deacetylation (PubMed:15226430). In addition to protein deacetylase activity, also has protein-lysine deacylase activity: acts as a protein decrotonylase by mediating decrotonylation ((2E)-butenoyl) of histones (PubMed:30279482).

{ECO:0000250|UniProtKB:Q13547, ECO:0000269|PubMed:10615135, ECO:0000269|PubMed:15226430, ECO:0000269|PubMed:15542849, ECO:0000269|PubMed:17707228, ECO:0000269|PubMed:21960634, ECO:0000269|PubMed:24736997, ECO:0000269|PubMed:30279482}.

Molecular Weight: 55.1 kDa

UniProt: [O09106](#)

Pathways: [Neurotrophin Signaling Pathway](#), [Intracellular Steroid Hormone Receptor Signaling Pathway](#), [Regulation of Intracellular Steroid Hormone Receptor Signaling](#), [Mitotic G1-G1/S Phases](#), [Regulation of Muscle Cell Differentiation](#), [Skeletal Muscle Fiber Development](#), [Negative Regulation of intrinsic apoptotic Signaling](#), [Embryonic Body Morphogenesis](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months