

Datasheet for ABIN7562295

CHEK1 Protein (AA 1-476) (His tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	CHEK1
Protein Characteristics:	AA 1-476
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CHEK1 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Purpose:	Custom-made recombinat Chek1 Protein expressed in mammalian cells.
Sequence:	<p>MAVPFVEDWD LVQTLGEGAY GEVQLAVNRI TEEAVAVKIV DMKRAIDCPE NIKKEICINK MLSHENVVKF YGHRREGHIQ YLFLEYCSGG ELFDRIEPI GMPEQDAQRF FHQLMAGVVY LHGIGITHRD IKPENLLLDE RDNLKISDFG LATVFRHNNR ERLNKMCGT LPYVAPELLK RKEFHAEPVD VWSCGIVLTA MLAGELPWDQ PSDSCQEYSD WKEKKTYPNP WKKIDSAPLA LLHKILVETP SARITIPDIK KDRWYNKPLN RGAKRPRATS GGMSESSSGF SKHIHSNLDF SPVNNGSSEE TVKFSSSQPE PRTGLSLWDT GPSNVDKLVQ GISFSQPTCP EHMLVNSQLL GTPGSSQNPW QRLVKRMTRF FTKLDADKSY QCLKETFEKL GYQWKKSCMN QVTVSTTDRR NNKLIFKINL VEMDEKILVD FRLSKGDGLE FKRHFLKIKG KLSDVSSQK VWFPVT Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.</p>

Product Details

Characteristics:

Key Benefits:

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

Grade:

custom-made

Target Details

Target:

CHEK1

Alternative Name:

Chk1 ([CHEK1 Products](#))

Background:

Serine/threonine-protein kinase Chk1 (EC 2.7.11.1) (CHK1 checkpoint homolog) (Checkpoint kinase-1),FUNCTION: Serine/threonine-protein kinase which is required for checkpoint-mediated cell cycle arrest and activation of DNA repair in response to the presence of DNA damage or unreplicated DNA (PubMed:10859163, PubMed:10859164, PubMed:15261141). May also negatively regulate cell cycle progression during unperturbed cell cycles (PubMed:10859163, PubMed:10859164, PubMed:15261141). This regulation is achieved by a number of mechanisms that together help to preserve the integrity of the genome (PubMed:10859163, PubMed:10859164, PubMed:15261141). Recognizes the substrate consensus sequence [R-X-X-S/T] (PubMed:10859163, PubMed:10859164, PubMed:15261141). Binds to and phosphorylates CDC25A, CDC25B and CDC25C. Phosphorylation of CDC25A at 'Ser-178' and 'Thr-507' and phosphorylation of CDC25C at 'Ser-216' creates binding sites for 14-3-3 proteins which inhibit CDC25A and CDC25C. Phosphorylation of CDC25A at 'Ser-76', 'Ser-124', 'Ser-178', 'Ser-279' and 'Ser-293' promotes proteolysis of CDC25A. Phosphorylation of

Target Details

CDC25A at 'Ser-76' primes the protein for subsequent phosphorylation at 'Ser-79', 'Ser-82' and 'Ser-88' by NEK11, which is required for polyubiquitination and degradation of CDC25A. Inhibition of CDC25 leads to increased inhibitory tyrosine phosphorylation of CDK-cyclin complexes and blocks cell cycle progression. Also phosphorylates NEK6. Binds to and phosphorylates RAD51 at 'Thr-309', which promotes the release of RAD51 from BRCA2 and enhances the association of RAD51 with chromatin, thereby promoting DNA repair by homologous recombination. Phosphorylates multiple sites within the C-terminus of TP53, which promotes activation of TP53 by acetylation and promotes cell cycle arrest and suppression of cellular proliferation. Also promotes repair of DNA cross-links through phosphorylation of FANCE. Binds to and phosphorylates TLK1 at 'Ser-743', which prevents the TLK1-dependent phosphorylation of the chromatin assembly factor ASF1A. This may enhance chromatin assembly both in the presence or absence of DNA damage. May also play a role in replication fork maintenance through regulation of PCNA (By similarity). May regulate the transcription of genes that regulate cell-cycle progression through the phosphorylation of histones. Phosphorylates histone H3.1 (to form H3T11ph), which leads to epigenetic inhibition of a subset of genes (PubMed:18243098). May also phosphorylate RB1 to promote its interaction with the E2F family of transcription factors and subsequent cell cycle arrest. Phosphorylates SPRTN, promoting SPRTN recruitment to chromatin (By similarity). Reduces replication stress and activates the G2/M checkpoint, by phosphorylating and inactivating PABIR1/FAM122A and promoting the serine/threonine-protein phosphatase 2A-mediated dephosphorylation and stabilization of WEE1 levels and activity (By similarity). {ECO:0000250|UniProtKB:O14757, ECO:0000269|PubMed:10859163, ECO:0000269|PubMed:10859164, ECO:0000269|PubMed:15261141, ECO:0000269|PubMed:18243098}.

Molecular Weight: 54.4 kDa

UniProt: [O35280](#)

Pathways: [p53 Signaling](#), [Apoptosis](#), [Cell Division Cycle](#), [DNA Damage Repair](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months