

Datasheet for ABIN7562299

NMI Protein (AA 1-314) (His tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	NMI
Protein Characteristics:	AA 1-314
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This NMI protein is labelled with His tag.

Product Details

Purpose:	Custom-made recombinant Nmi Protein expressed in mammalian cells.
Sequence:	<p>MDADKDNKIQ ACDERSAEMD DMRGEQSMGL VHEIMSENKE LDEEIKKLEA ELQSDAREFQ IKENVPEKKL KLTSVESPKD GCHFSNSSCS FQVSSQILYE LQEGQALITF EKEEVAQNV SMGNHVVQME GTPVKVSAHP VPLNTGVRFQ VHVDISKMKI NVTGIPDELS EEQTRDKLEL SFCKSRNGGG EVESVDYDRK SRSAVITFVE TGVVDKILKK KTYPLYMNQK CHSVAVSPCI ERCLEKYQVF SAVSKKTVLL TGLEGIPVDE ETGEDLLNIH FQRKNNGGGE VEVVKCSLDQ SFAAYFKEEA RETI Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.</p>
Specificity:	If you are looking for a specific domain and are interested in a partial protein or a different isoform, please contact us regarding an individual offer.
Characteristics:	Key Benefits:

Product Details

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:	> 90 % as determined by Bis-Tris PAGE, anti-tag ELISA, Western Blot and analytical SEC (HPLC)
Grade:	custom-made

Target Details

Target:	NMI
Alternative Name:	Nmi (NMI Products)
Background:	N-myc-interactor (Nmi) (N-myc and STAT interactor),FUNCTION: Acts as a signaling pathway regulator involved in innate immune system response (PubMed:23956435). In response to interleukin 2/IL2 and interferon IFN-gamma/IFNG, interacts with signal transducer and activator of transcription/STAT which activate the transcription of downstream genes involved in a multitude of signals for development and homeostasis (By similarity). Enhances the recruitment of CBP/p300 coactivators to STAT1 and STAT5, resulting in increased STAT1- and STAT5-dependent transcription (By similarity). In response to interferon IFN-alpha, associates in a complex with transcriptional regulator IFI35 to regulate immune response, the complex formation prevents proteasome-mediated degradation of IFI35 (By similarity). In complex with IFI35, negatively regulates nuclear factor NF-kappa-B signaling by inhibiting the nuclear translocation, activation and transcription of NF-kappa-B subunit p65/RELA, resulting in the inhibition of endothelial cell proliferation, migration and re-endothelialization of injured arteries (By similarity). Negatively regulates virus-triggered type I interferon/IFN production by inducing proteasome-dependent degradation of IRF7, a transcriptional regulator of type I IFN, thereby

Target Details

interfering with cellular antiviral responses (PubMed:23956435). Beside its role as an intracellular signaling pathway regulator, also functions extracellularly as damage-associated molecular patterns (DAMPs) to promote inflammation, when actively released by macrophage to the extracellular space during cell injury or pathogen invasion (By similarity). Macrophage-secreted NMI activates NF-kappa-B signaling in adjacent macrophages through Toll-like receptor 4/TLR4 binding and activation, thereby inducing NF-kappa-B translocation from the cytoplasm into the nucleus which promotes the release of pro-inflammatory cytokines (By similarity). {ECO:0000250|UniProtKB:Q13287, ECO:0000269|PubMed:23956435}.

Molecular Weight: 35.2 kDa

UniProt: [O35309](#)

Application Details

Application Notes: We expect the protein to work for functional studies. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: 12 months