

Datasheet for ABIN7562504

## Aurora Kinase B Protein (AURKB) (AA 1-345) (His tag)



[Go to Product page](#)

### Overview

Quantity:	1 mg
Target:	Aurora Kinase B (AURKB)
Protein Characteristics:	AA 1-345
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Aurora Kinase B protein is labelled with His tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB)

### Product Details

Purpose:	Custom-made recombinat Aurkb Protein expressed in mammalien cells.
Sequence:	<p>MAQKENAYPW PYGSKTSQSG LNTLSQRVLR KEPATTSALA LVNRFNSQST AAPGQKLAEN</p> <p>KSQGSTASQG SQNKQPFTID NFEIGRPLGK GKFGNVYLAR EKKSRLFVAL KILFKSQIEK</p> <p>EGVEHQLRRE IEIQAHLKHP NILQLYNYFY DQQRIYLILE YAPRGELYKE LQKSRTFDEQ</p> <p>RTATIMEELS DALTYCHKKK VIHRDIKPEN LLLGLQGELK IADFGWSVHA PSLRRKTMCG</p> <p>TLDYLPPEMI EGRMHNMVD LWCIGVLCYE LMVGNPPFES PSHSETYRRI VKVDLKFPSS</p> <p>VPSGAQDLIS KLLKHNPWQR LPLAEVAAHP WVRANSRRVL PPSAL <b>Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.</b></p>
Characteristics:	Key Benefits:

## Product Details

---

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

---

Purity:	> 90 % as determined by Bis-Tris Page, Western Blot
---------	---

---

Grade:	custom-made
--------	-------------

---

## Target Details

---

Target:	Aurora Kinase B (AURKB)
---------	-------------------------

---

Alternative Name:	Aurkb ( <a href="#">AURKB Products</a> )
-------------------	--

---

Background:	<p>Aurora kinase B (EC 2.7.11.1) (Aurora 1) (Aurora- and IPL1-like midbody-associated protein 1) (Aurora/IPL1-related kinase 2) (ARK-2) (Aurora-related kinase 2) (STK-1) (Serine/threonine-protein kinase 12) (Serine/threonine-protein kinase 5) (Serine/threonine-protein kinase aurora-B),FUNCTION: Serine/threonine-protein kinase component of the chromosomal passenger complex (CPC), a complex that acts as a key regulator of mitosis (By similarity). The CPC complex has essential functions at the centromere in ensuring correct chromosome alignment and segregation and is required for chromatin-induced microtubule stabilization and spindle assembly (By similarity). Involved in the bipolar attachment of spindle microtubules to kinetochores and is a key regulator for the onset of cytokinesis during mitosis (By similarity). Required for central/midzone spindle assembly and cleavage furrow formation (By similarity). Key component of the cytokinesis checkpoint, a process required to delay abscission to prevent both premature resolution of intercellular chromosome bridges and accumulation of DNA damage: phosphorylates CHMP4C, leading to retain abscission-competent VPS4 (VPS4A and/or VPS4B) at the midbody ring until abscission checkpoint signaling is terminated at late</p>
-------------	--

---

Target Details

cytokinesis (By similarity). AURKB phosphorylates the CPC complex subunits BIRC5/survivin, CDCA8/borealin and INCENP (By similarity). Phosphorylation of INCENP leads to increased AURKB activity (By similarity). Other known AURKB substrates involved in centromeric functions and mitosis are CENPA, DES/desmin, GPAF, KIF2C, NSUN2, RACGAP1, SEPTIN1, VIM/vimentin, HASPIN, and histone H3 (By similarity). A positive feedback loop involving HASPIN and AURKB contributes to localization of CPC to centromeres (By similarity). Phosphorylation of VIM controls vimentin filament segregation in cytokinetic process, whereas histone H3 is phosphorylated at 'Ser-10' and 'Ser-28' during mitosis (H3S10ph and H3S28ph, respectively) (PubMed:11784863). AURKB is also required for kinetochore localization of BUB1 and SGO1 (By similarity). Phosphorylation of p53/TP53 negatively regulates its transcriptional activity (By similarity). Key regulator of active promoters in resting B- and T-lymphocytes: acts by mediating phosphorylation of H3S28ph at active promoters in resting B-cells, inhibiting RNF2/RING1B-mediated ubiquitination of histone H2A and enhancing binding and activity of the USP16 deubiquitinase at transcribed genes (PubMed:24034696). Acts as an inhibitor of CGAS during mitosis: catalyzes phosphorylation of the N-terminus of CGAS during the G2-M transition, blocking CGAS liquid phase separation and activation, and thereby preventing CGAS-induced autoimmunity (By similarity). Phosphorylates KRT5 during anaphase and telophase (PubMed:29518391). {ECO:0000250|UniProtKB:Q96GD4, ECO:0000269|PubMed:11784863, ECO:0000269|PubMed:24034696, ECO:0000269|PubMed:29518391}.

Molecular Weight:	39.4 kDa
UniProt:	<a href="#">O70126</a>
Pathways:	<a href="#">TCR Signaling</a> , <a href="#">Cell Division Cycle</a> , <a href="#">Maintenance of Protein Location</a> , <a href="#">Hepatitis C</a> , <a href="#">Toll-Like Receptors Cascades</a>

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.

Handling

Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months