

Datasheet for ABIN7562610
IGF2BP1 Protein (AA 1-577) (His tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	IGF2BP1
Protein Characteristics:	AA 1-577
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This IGF2BP1 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Purpose:	Custom-made recombinat Igf2bp1 Protein expressed in mammalian cells.
Sequence:	<p>MNKLYIGNLN ESVTPADLEK VFAEHKISYS GQFLVKSGYA FVDCPDEHWA MKAIETFSGK VELQGKRLEI EHSVPPKKQRS RKIQIRNIPP QLRWEVLDSL LAQYGTVENC EQVNTESETA VVNVTSYNRE QTRQAIMKLN GHQLENHALK VSYIPDEQIT QGPENGRRGG FGSRGQPRQG SPVAAGAPAK QQPVDIPLRL LVPTQYVGAI IGKEGATIRN ITKQTQSKID VHRKENAGAA EKAISVHSTP EGCSSACKMI LEIMHKEAKD TKTADDEVPLK ILAHNNFVGR LIGKEGRNLK KVEQDTETKI TISSLQDLTL YNPERTITVK GAIENCCRAE QEIMKKVREA YENDVAAMSL QSHLIPGLNL AAVGLFPASS SAVPPPPSSV TGAAPYSSFM QAPEQEMVQV FIPAQAVGAI IGKKGQHIKQ LSRFASASIK IAPPETPDSK VRMVVITGPP EAQFKAQGRI YGKLKEENFF GPKEEVKLET HIRVPASAAG RVIGKGGKTV NELQNLTAEE VVPRDQTPD ENDQVIVKII GHFYASQMAQ RKIRDILAQV KQQHQKGQSN LAQARRK Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity</p>

of the protein could make another tag necessary. In case you have a special request, please contact us.

Characteristics:	<p>Key Benefits:</p> <ul style="list-style-type: none">• Made to order protein - from design to production - by highly experienced protein experts.• Protein expressed in mammalian cells and purified in one-step affinity chromatography• The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.• State-of-the-art algorithm used for plasmid design (Gene synthesis). <p>This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.</p> <p>If you are not interested in a full length protein, please contact us for individual protein fragments.</p> <p>The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.</p>
Purity:	> 90 % as determined by Bis-Tris Page, Western Blot
Grade:	custom-made

Target Details

Target:	IGF2BP1
Alternative Name:	Igf2bp1 (IGF2BP1 Products)
Background:	<p>Insulin-like growth factor 2 mRNA-binding protein 1 (IGF2 mRNA-binding protein 1) (IMP-1) (Coding region determinant-binding protein) (CRD-BP) (IGF-II mRNA-binding protein 1) (VICKZ family member 1) (Zipcode-binding protein 1) (ZBP-1),FUNCTION: RNA-binding factor that recruits target transcripts to cytoplasmic protein-RNA complexes (mRNPs). This transcript 'caging' into mRNPs allows mRNA transport and transient storage. It also modulates the rate and location at which target transcripts encounter the translational apparatus and shields them from endonuclease attacks or microRNA-mediated degradation. Preferentially binds to N6-methyladenosine (m6A)-containing mRNAs and increases their stability (By similarity). Regulates localized beta-actin/ACTB mRNA translation, a crucial process for cell polarity, cell migration and neurite outgrowth. Co-transcriptionally associates with the ACTB mRNA in the nucleus. This binding involves a conserved 54-nucleotide element in the ACTB mRNA 3'-UTR,</p>

Target Details

known as the 'zipcode'. The RNP thus formed is exported to the cytoplasm, binds to a motor protein and is transported along the cytoskeleton to the cell periphery. During transport, prevents ACTB mRNA from being translated into protein. When the RNP complex reaches its destination near the plasma membrane, IGF2BP1 is phosphorylated. This releases the mRNA, allowing ribosomal 40S and 60S subunits to assemble and initiate ACTB protein synthesis. Monomeric ACTB then assembles into the subcortical actin cytoskeleton (By similarity). During neuronal development, key regulator of neurite outgrowth, growth cone guidance and neuronal cell migration, presumably through the spatiotemporal fine tuning of protein synthesis, such as that of ACTB (By similarity). May regulate mRNA transport to activated synapses (By similarity). Binds to the 3'-UTR of CD44 mRNA and stabilizes it, hence promotes cell adhesion and invadopodia formation in cancer cells (By similarity). Binds to the oncofetal H19 transcript and regulates its localization (By similarity). Binds to and stabilizes BTRC/FBW1A mRNA (By similarity). Binds to the adenine-rich autoregulatory sequence (ARS) located in PABPC1 mRNA and represses its translation. PABPC1 mRNA-binding is stimulated by PABPC1 protein. Prevents BTRC/FBW1A mRNA degradation by disrupting microRNA-dependent interaction with AGO2 (By similarity). During cellular stress, such as oxidative stress or heat shock, stabilizes target mRNAs that are recruited to stress granules, including CD44, IGF2, MAPK4, MYC, PTEN, RAPGEF2 and RPS6KA5 transcripts (By similarity). Interacts with GAP43 transcript and transports it to axons. Binds to the 3'-UTR of IGF2 mRNA by a mechanism of cooperative and sequential dimerization and regulates IGF2 mRNA subcellular localization and translation. Binds to MYC mRNA, in the coding region instability determinant (CRD) of the open reading frame (ORF), hence prevents MYC cleavage by endonucleases and possibly microRNA targeting to MYC-CRD. Binding to MYC mRNA is enhanced by m6A-modification of the CRD (By similarity). Binds to and stabilizes ABCB1/MDR-1 mRNA. Binds to the neuron-specific TAU mRNA and regulates its localization. Plays a direct role in the transport and translation of transcripts required for axonal regeneration in adult sensory neurons. During interstitial wound repair, interacts with and stabilizes PTGS2 transcript. PTGS2 mRNA stabilization may be crucial for colonic mucosal wound healing. {ECO:0000250, ECO:0000250|UniProtKB:Q9NZI8, ECO:0000269|PubMed:15355996, ECO:0000269|PubMed:17264115, ECO:0000269|PubMed:21964071, ECO:0000269|PubMed:22465430}.

Molecular Weight:	63.5 kDa
UniProt:	O88477

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
--------------------	---

Application Details

as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: 12 months