

Datasheet for ABIN7562737

Mitochondrially Encoded ATP Synthase 8 (MT-ATP8) (AA 1-67) protein (Fc Tag)



Go to Product page

Overview	vervi	UV	۷
----------	-------	----	---

Quantity:	1 mg
Target:	Mitochondrially Encoded ATP Synthase 8 (MT-ATP8)
Protein Characteristics:	AA 1-67
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	Fc Tag
Application:	Western Blotting (WB), SDS-PAGE (SDS)
Product Details	
Purpose:	Custom-made recombinat Mtatp8 Protein expressed in mammalien cells.
Sequence:	MPQLDTSTWF ITIISSMITL FILFQLKVSS QTFPLAPSPK SLTTMKVKTP WELKWTKIYL PHSLPQQ
	Sequence without tag. The proposed Purification-Tag is based on experiences with the
	expression system, a different complexity of the protein could make another tag necessary.
	In case you have a special request, please contact us.
Characteristics:	Key Benefits:
	Made to order protein - from design to production - by highly experienced protein experts.
	Protein expressed in mammalien cells and purified in one-step affinity chromatography
	The optimized expression system ensures reliability for intracellular, secreted and
	transmembrane proteins.
	State-of-the-art algorithm used for plasmid design (Gene synthesis).

Restrictions:

Product Details	
	experts in the lab try to ensure that you receive soluble protein.
	If you are not interested in a full length protein, please contact us for individual protein fragments.
	The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.
Purity:	> 90 % as determined by Bis-Tris Page, Western Blot
Grade:	custom-made
Target Details	
Target:	Mitochondrially Encoded ATP Synthase 8 (MT-ATP8)
Alternative Name:	Mtatp8 (MT-ATP8 Products)
Background:	ATP synthase protein 8 (A6L) (F-ATPase subunit 8),FUNCTION: Mitochondrial membrane ATP synthase (F(1)F(0) ATP synthase or Complex V) produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F(1) - containing the extramembraneous catalytic core and F(0) - containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F(1) is coupled via a rotary mechanism of the central stalk subunits to proton translocation. Part of the complex F(0) domain. Minor subunit located with subunit a in the membrane (By similarity). {EC0:0000250}.
Molecular Weight:	7.8 kDa
UniProt:	P03930
Pathways:	Proton Transport, Ribonucleoside Biosynthetic Process
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Doctrictions:	For Doggarch Llog only

For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months