

[Go to Product page](#)

Datasheet for ABIN7562810

ALKBH1 Protein (AA 1-389) (His tag)

Overview

Quantity:	1 mg
Target:	ALKBH1
Protein Characteristics:	AA 1-389
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This ALKBH1 protein is labelled with His tag.

Product Details

Purpose:	Custom-made recombinant Alkbh1 Protein expressed in mammalian cells.
Sequence:	MGKMAAVAS LATLAAEPRE DAFRKLFRFY RQSRPGTADL GAVIDFSEAH LARSPKPGVP QVVRFP LNVS SVTERDAERV GLEPVSKWRA YGLEGYPGFI FIPNPFLPGC QRHWVKQCLK LYSQPNVCN LDKHMTKEET QGLWEQSKEV LRSKEVTKRR PRSLLERLRW VTLGYHYNWD SKKYSADHYT PFPSDLAFLS EQVATACGFQ GFQAEAGILN YYRLDSTLGI HVDRSELDHS KPLLSFSFGQ SAIFLLGGLK RDEAPTAMFM HSGDIMVMSG FSRLLNHAVP RVLPHPDGEC LPHCLETPLP AVLPSNSLVE PCSVEDWQVC ATYLRTARVN MTVRQVLATG QDFPLEPVEE TKRDIAADGL CHLHDPNSPV KRKRLNPNS Sequence without tag. The proposed Purification- Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.
Specificity:	If you are looking for a specific domain and are interested in a partial protein or a different isoform, please contact us regarding an individual offer.

Product Details

Characteristics:

Key Benefits:

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris PAGE, anti-tag ELISA, Western Blot and analytical SEC (HPLC)

Grade:

custom-made

Target Details

Target:

ALKBH1

Alternative Name:

Alkbh1 ([ALKBH1 Products](#))

Background:

Nucleic acid dioxygenase ALKBH1 (EC 1.14.11.-) (Alkylated DNA repair protein alkB homolog 1) (Alpha-ketoglutarate-dependent dioxygenase ABH1) (DNA 6mA demethylase) (DNA N6-methyl adenine demethylase ALKBH1) (EC 1.14.11.51) (DNA lyase ABH1) (EC 4.2.99.18) (DNA oxidative demethylase ALKBH1) (EC 1.14.11.33) (mRNA N(3)-methylcytidine demethylase) (EC 1.14.11.-) (tRNA N1-methyl adenine demethylase) (EC 1.14.11.-),FUNCTION: Dioxygenase that acts as on nucleic acids, such as DNA and tRNA (PubMed:27027282, PubMed:27745969). Requires molecular oxygen, alpha-ketoglutarate and iron (PubMed:27027282). A number of activities have been described for this dioxygenase, but recent results suggest that it mainly acts as on tRNAs and mediates their demethylation or oxidation depending on the context and subcellular compartment (By similarity). Mainly acts as a tRNA demethylase by removing N(1)-methyladenine from various tRNAs, with a preference for N(1)-methyladenine at position 58 (m1A58) present on a stem loop structure of tRNAs (PubMed:27745969). Acts as a regulator of translation initiation and elongation in response to glucose deprivation: regulates both

Target Details

translation initiation, by mediating demethylation of tRNA(Met), and translation elongation, N(1)-methyladenine-containing tRNAs being preferentially recruited to polysomes to promote translation elongation (By similarity). In mitochondrion, specifically interacts with mt-tRNA(Met) and mediates oxidation of mt-tRNA(Met) methylated at cytosine(34) to form 5-formylcytosine (f(5)c) at this position (By similarity). mt-tRNA(Met) containing the f(5)c modification at the wobble position enables recognition of the AUA codon in addition to the AUG codon, expanding codon recognition in mitochondrial translation (By similarity). Specifically demethylates DNA methylated on the 6th position of adenine (N(6)-methyladenosine) DNA (PubMed:27027282). N(6)-methyladenosine (m6A) DNA is present at some L1 elements in embryonic stem cells and probably promotes their silencing (PubMed:27027282). Demethylates mRNAs containing N(3)-methylcytidine modification (By similarity). Also able to repair alkylated single-stranded DNA by oxidative demethylation, but with low activity (By similarity). Also has DNA lyase activity and introduces double-stranded breaks at abasic sites: cleaves both single-stranded DNA and double-stranded DNA at abasic sites, with the greatest activity towards double-stranded DNA with two abasic sites (By similarity). DNA lyase activity does not require alpha-ketoglutarate and iron and leads to the formation of an irreversible covalent protein-DNA adduct with the 5' DNA product (By similarity). DNA lyase activity is not required during base excision repair and class switch recombination of the immunoglobulin heavy chain during B lymphocyte activation (PubMed:23825659). May play a role in placental trophoblast lineage differentiation (PubMed:18163532). {ECO:0000250|UniProtKB:Q13686, ECO:0000269|PubMed:18163532, ECO:0000269|PubMed:23825659, ECO:0000269|PubMed:27027282, ECO:0000269|PubMed:27745969}.

Molecular Weight: 43.7 kDa

UniProt: [P0CB42](#)

Pathways: [DNA Damage Repair](#)

Application Details

Application Notes: We expect the protein to work for functional studies. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions: For Research Use only

Handling

Format: Liquid

Handling

Buffer: The buffer composition is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: 12 months