

Datasheet for ABIN7562945

ID1 Protein (AA 1-168) (His tag)



_					
	W	0	rv	10	W

Quantity:	1 mg
Target:	ID1
Protein Characteristics:	AA 1-168
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This ID1 protein is labelled with His tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB)
Product Details	

Product Details	
Purpose:	Custom-made recombinat Id1 Protein expressed in mammalien cells.
Sequence:	MKVASGSAAA AAGPSCSLKA GRTAGEVVLG LSEQSVAISR CAGTRLPALL DEQQVNVLLY
	DMNGCYSRLK ELVPTLPQNR KVSKVEILQH VIDYIRDLQL ELNSESEVGT TGGRGLPVRA
	PLSTLNGEIS ALAAEVRSES EYYIILQWET EATGGGCPPS LLFRRIAI Sequence without tag. The
	proposed Purification-Tag is based on experiences with the expression system, a different
	complexity of the protein could make another tag necessary. In case you have a special
	request, please contact us.
Characteristics:	Key Benefits:
	Made to order protein - from design to production - by highly experienced protein experts.
	 Protein expressed in mammalien cells and purified in one-step affinity chromatography
	 The optimized expression system ensures reliability for intracellular, secreted and
	transmembrane proteins.

• State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

Grade:

custom-made

Target Details

Alternative Name:

Target:	ID1

Id1 (ID1 Products)

Background:

DNA-binding protein inhibitor ID-1 (Inhibitor of DNA binding 1) (Inhibitor of differentiation 1),FUNCTION: Transcriptional regulator (lacking a basic DNA binding domain) which negatively regulates the basic helix-loop-helix (bHLH) transcription factors by forming heterodimers and inhibiting their DNA binding and transcriptional activity. Implicated in regulating a variety of cellular processes, including cellular growth, senescence, differentiation, apoptosis, angiogenesis, and neoplastic transformation. Inhibits skeletal muscle and cardiac myocyte differentiation. Regulates the circadian clock by repressing the transcriptional activator activity of the CLOCK-BMAL1 heterodimer. {ECO:0000269|PubMed:16556596,

Molecular Weight:

17.9 kDa

UniProt:

P20067

Application Details

Application Notes:

In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Application Details

Storage Comment:

Expiry Date:

Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	The buffer composition is at the discretion of the manufacturer.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	

Store at -80°C.

12 months