

Datasheet for ABIN7562993

## ZFP36L1 Protein (AA 1-338) (His tag)



[Go to Product page](#)

### Overview

Quantity:	1 mg
Target:	ZFP36L1
Protein Characteristics:	AA 1-338
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This ZFP36L1 protein is labelled with His tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB)

### Product Details

Purpose:	Custom-made recombinat Zfp36l1 Protein expressed in mammalian cells.
Sequence:	<p>MTTTLVSATI FDLSEVLCKG NKMLNYSTPS AGGCLLDRKA VGTPAGGGFP RRHSVTLPS</p> <p>KFHQNQLLSS LKGEPAPSLs SRDSRFRDRS FSEGGERLLP TQKQPGSGQV NSSRYKTELC</p> <p>RPFEENGACK YGDKCQFAHG IHELRLSLTRH PKYKTELCRT FHTIGFCPYG PRCHFIHNAE</p> <p>ERRALAGGRD LSADRPRLQH SFSFAGFPSA AATAAATGLL DSPTSITPPP ILSADDLLGS</p> <p>PTLPDGTNNP FAFSSQELAS LFAPSMGLPG GGSPTTFLFR PMSESPHMFDP SPPSPQDSLS</p> <p>DHEGYLSSSS SSHSGSDSPT LDNSRRLPIF SRLSISDD <b>Sequence without tag. The proposed</b></p> <p><b>Purification-Tag is based on experiences with the expression system, a different complexity</b></p> <p><b>of the protein could make another tag necessary. In case you have a special request, please</b></p> <p><b>contact us.</b></p>
Characteristics:	Key Benefits:

## Product Details

---

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

---

Purity:	> 90 % as determined by Bis-Tris Page, Western Blot
---------	---

---

Grade:	custom-made
--------	-------------

---

## Target Details

---

Target:	ZFP36L1
---------	---------

---

Alternative Name:	Zfp36l1 ( <a href="#">ZFP36L1 Products</a> )
-------------------	--

---

Background:	<p>MRNA decay activator protein ZFP36L1 (Butyrate response factor 1) (TPA-induced sequence 11b) (Zinc finger protein 36, C3H1 type-like 1) (ZFP36-like 1),FUNCTION: Zinc-finger RNA-binding protein that destabilizes several cytoplasmic AU-rich element (ARE)-containing mRNA transcripts by promoting their poly(A) tail removal or deadenylation, and hence provide a mechanism for attenuating protein synthesis (PubMed:22701344, PubMed:24700863, PubMed:24733888, PubMed:27102483). Acts as a 3'-untranslated region (UTR) ARE mRNA-binding adapter protein to communicate signaling events to the mRNA decay machinery (By similarity). Functions by recruiting the CCR4-NOT deadenylating complex and components of the cytoplasmic RNA decay machinery to the bound ARE-containing mRNAs, and hence promotes ARE-mediated mRNA deadenylation and decay processes (By similarity). Induces also the degradation of ARE-containing mRNAs even in absence of poly(A) tail (By similarity). Binds to 3'-UTR ARE of numerous mRNAs (PubMed:22701344, PubMed:24700863, PubMed:24733888). Positively regulates early adipogenesis by promoting ARE-mediated mRNA decay of immediate early genes (IEGs) (PubMed:22701344). Promotes ARE-mediated mRNA</p>
-------------	--

---

decay of mineralocorticoid receptor NR3C2 mRNA in response to hypertonic stress (PubMed:24700863). Negatively regulates hematopoietic/erythroid cell differentiation by promoting ARE-mediated mRNA decay of the transcription factor STAT5B mRNA (By similarity). Positively regulates monocyte/macrophage cell differentiation by promoting ARE-mediated mRNA decay of the cyclin-dependent kinase CDK6 mRNA (By similarity). Promotes degradation of ARE-containing pluripotency-associated mRNAs in embryonic stem cells (ESCs), such as NANOG, through a fibroblast growth factor (FGF)-induced MAPK-dependent signaling pathway, and hence attenuates ESC self-renewal and positively regulates mesendoderm differentiation (PubMed:24733888). May play a role in mediating pro-apoptotic effects in malignant B-cells by promoting ARE-mediated mRNA decay of BCL2 mRNA (By similarity). In association with ZFP36L2 maintains quiescence on developing B lymphocytes by promoting ARE-mediated decay of several mRNAs encoding cell cycle regulators that help B cells progress through the cell cycle, and hence ensuring accurate variable-diversity-joining (VDJ) recombination and functional immune cell formation (PubMed:27102483). Together with ZFP36L2 is also necessary for thymocyte development and prevention of T-cell acute lymphoblastic leukemia (T-ALL) transformation by promoting ARE-mediated mRNA decay of the oncogenic transcription factor NOTCH1 mRNA (PubMed:20622884). Involved in the delivery of target ARE-mRNAs to processing bodies (PBs) (By similarity). In addition to its cytosolic mRNA-decay function, plays a role in the regulation of nuclear mRNA 3'-end processing, modulates mRNA 3'-end maturation efficiency of the DLL4 mRNA through binding with an ARE embedded in a weak noncanonical polyadenylation (poly(A)) signal in endothelial cells (By similarity). Also involved in the regulation of stress granule (SG) and P-body (PB) formation and fusion (By similarity). Plays a role in vasculogenesis and endocardial development (PubMed:15226444, PubMed:17013884). Involved in the regulation of keratinocyte proliferation, differentiation and apoptosis (By similarity). Plays a role in myoblast cell differentiation (PubMed:17889962). {ECO:0000250|UniProtKB:P17431, ECO:0000250|UniProtKB:Q07352, ECO:0000269|PubMed:15226444, ECO:0000269|PubMed:17013884, ECO:0000269|PubMed:17889962, ECO:0000269|PubMed:20622884, ECO:0000269|PubMed:22701344, ECO:0000269|PubMed:24700863, ECO:0000269|PubMed:24733888, ECO:0000269|PubMed:27102483}.

Molecular Weight: 36.4 kDa

UniProt: [P23950](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies

Application Details

as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: 12 months