

Datasheet for ABIN7563068

## DPP4 Protein (AA 1-760) (His tag)



[Go to Product page](#)

### Overview

Quantity:	1 mg
Target:	DPP4
Protein Characteristics:	AA 1-760
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This DPP4 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

### Product Details

Purpose:	Custom-made recombinat Dpp4 Protein expressed in mammalian cells.
Sequence:	<p>MKTPWKVLLG LLGVAALVTI ITVPIVLLSK DEAAADSRRT YSLADYLKST FRVKSYSLLWW</p> <p>VSDFEYLYKQ ENNILLNNAE HGNSSIFLEN STFESFGYHS VSPDRFLVLL EYNYVKQWRH</p> <p>SYTASYNIYD VNKRQLITEE KIPNNTQWIT WSPGHLKLAY VWKNDIYVKV EPHLPShRIT</p> <p>STGEENVIYN GITDWVYEEE VFGAYSALWW SPNNTFLAYA QFNDTGVPLI EYSFYSDSL</p> <p>QYPKTVWIPY PKAGAVNPTV KFFIVNIDSL SSSSSAAPIQ IPAPASVARG DHYLCDVVWA</p> <p>TEERISLQWL RRIQNYSVMA ICDYDKINLT WNCPSQQHV EMSTTGWVGR FRPAEPHFTS</p> <p>DGSSFYKIIS DKDGYKHICH FPKDKKDCF ITKGAWAVIS IEALTSYDLY YISNQYKEMP</p> <p>GGRNLYKIQL TDHTNVKCLS CDLNPERCQY YAVSFSKEAK YYQLGCWGPGLPLYTLHRST</p> <p>DHKELRVLED NSALDRMLQD VQMPSKKLDF IVLNETRFWY QMILPPHFDK SKKYPLLLDV</p> <p>YAGPCSQKAD ASFRLNWATY LASTENIIVA SFDGRGSGYQ GDKIMHAINR RLGTLVEDQ</p> <p>IEAARQFVKM GFVDSKRVAI WGWSYGGYVT SMVLGSGSGV FKCGIAPAPV SRWEYYDSVY</p>

Product Details

TERYMGLPIP EDNLDHYRNS TVMSRAEHFK QVEYLLIHGT ADDNVHFQQS AQISKALVDA  
GVDFQAMWYT DEDHGIASST AHQHIYSHMS HFLQQCFSLH **Sequence without tag. The proposed Purification-Tag is based on experiences with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.**

Characteristics:	<div>Key Benefits:</div> <ul style="list-style-type: none"><li>• Made to order protein - from design to production - by highly experienced protein experts.</li><li>• Protein expressed in mammalian cells and purified in one-step affinity chromatography</li><li>• The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.</li><li>• State-of-the-art algorithm used for plasmid design (Gene synthesis).</li></ul> <p>This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.</p> <p>If you are not interested in a full length protein, please contact us for individual protein fragments.</p> <p>The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.</p>
------------------	---

Purity:	> 90 % as determined by Bis-Tris Page, Western Blot
---------	---

Grade:	custom-made
--------	-------------

Target Details

Target:	DPP4
Alternative Name:	Dpp4 (DPP4 Products)
Background:	Dipeptidyl peptidase 4 (EC 3.4.14.5) (Dipeptidyl peptidase IV) (DPP IV) (T-cell activation antigen CD26) (Thymocyte-activating molecule) (THAM) (CD antigen CD26) [Cleaved into: Dipeptidyl peptidase 4 membrane form (Dipeptidyl peptidase IV membrane form), Dipeptidyl peptidase 4 soluble form (Dipeptidyl peptidase IV soluble form)],FUNCTION: Cell surface glycoprotein receptor involved in the costimulatory signal essential for T-cell receptor (TCR)-mediated T-cell activation. Acts as a positive regulator of T-cell coactivation, by binding at least ADA, CAV1, IGF2R, and PTPRC. Its binding to CAV1 and CARD11 induces T-cell proliferation and NF-kappa-B activation in a T-cell receptor/CD3-dependent manner. Its interaction with ADA also regulates

## Target Details

lymphocyte-epithelial cell adhesion. In association with FAP is involved in the pericellular proteolysis of the extracellular matrix (ECM), the migration and invasion of endothelial cells into the ECM. May be involved in the promotion of lymphatic endothelial cells adhesion, migration and tube formation. When overexpressed, enhanced cell proliferation, a process inhibited by GPC3. Acts also as a serine exopeptidase with a dipeptidyl peptidase activity that regulates various physiological processes by cleaving peptides in the circulation, including many chemokines, mitogenic growth factors, neuropeptides and peptide hormones. Removes N-terminal dipeptides sequentially from polypeptides having unsubstituted N-termini provided that the penultimate residue is proline. {ECO:0000250|UniProtKB:P27487}.

Molecular Weight: 87.4 kDa

UniProt: [P28843](#)

Pathways: [Peptide Hormone Metabolism, Regulation of Leukocyte Mediated Immunity](#)

## Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: 12 months