

Datasheet for ABIN7563202 **HSF1 Protein (AA 1-525) (His tag)**



Overview

Quantity:	1 mg
Target:	HSF1
Protein Characteristics:	AA 1-525
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This HSF1 protein is labelled with His tag.

Product Details	
Purpose:	Custom-made recombinant Hsf1 Protein expressed in mammalian cells.
Sequence:	MDLAVGPGAA GPSNVPAFLT KLWTLVSDPD TDALICWSPS GNSFHVFDQG QFAKEVLPKY
	FKHNNMASFV RQLNMYGFRK VVHIEQGGLV KPERDDTEFQ HPCFLRGQEQ LLENIKRKVT
	SVSTLKSEDI KIRQDSVTRL LTDVQLMKGK QECMDSKLLA MKHENEALWR EVASLRQKHA
	QQQKVVNKLI QFLISLVQSN RILGVKRKIP LMLSDSNSAH SVPKYGRQYS LEHVHGPGPY
	SAPSPAYSSS SLYSSDAVTS SGPIISDITE LAPTSPLASP GRSIDERPLS SSTLVRVKQE
	PPSPPHSPRV LEASPGRPSS MDTPLSPTAF IDSILRESEP TPAASNTAPM DTTGAQAPAL
	PTPSTPEKCL SVACLDKNEL SDHLDAMDSN LDNLQTMLTS HGFSVDTSAL LDLFSPSVTM
	PDMSLPDLDS SLASIQELLS PQEPPRPIEA ENSNPDSGKQ LVHYTAQPLF LLDPDAVDTG
	SSELPVLFEL GESSYFSEGD DYTDDPTISL LTGTEPHKAK DPTVS Sequence without tag. The
	proposed Purification-Tag is based on experiences with the expression system, a different
	complexity of the protein could make another tag necessary. In case you have a special
	request, please contact us.

Product Details

Specificity: If you are looking for a specific domain and are interested in a partial protein or a different isoform, please contact us regarding an individual offer. Characteristics: Key Benefits: · Made to order protein - from design to production - by highly experienced protein experts. · Protein expressed in mammalian cells and purified in one-step affinity chromatography · The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins. • State-of-the-art algorithm used for plasmid design (Gene synthesis). This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein. If you are not interested in a full length protein, please contact us for individual protein fragments. The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified. Purity: > 90 % as determined by Bis-Tris PAGE, anti-tag ELISA, Western Blot and analytical SEC (HPLC) Grade: custom-made **Target Details** Target: HSF1 Alternative Name: Hsf1 (HSF1 Products) Background: Heat shock factor protein 1 (HSF 1) (Heat shock transcription factor 1) (HSTF 1),FUNCTION: Functions as a stress-inducible and DNA-binding transcription factor that plays a central role in the transcriptional activation of the heat shock response (HSR), leading to the expression of a large class of molecular chaperones, heat shock proteins (HSPs), that protect cells from cellular insult damage. In unstressed cells, is present in a HSP90-containing multichaperone complex

other stress stimuli, undergoes homotrimerization and activates HSP gene transcription through binding to site-specific heat shock elements (HSEs) present in the promoter regions of HSP genes. Upon heat shock stress, forms a chromatin-associated complex with TTC5/STRAP and p300/EP300 to stimulate HSR transcription, therefore increasing cell survival. Activation is reversible, and during the attenuation and recovery phase period of the HSR, returns to its

that maintains it in a non-DNA-binding inactivated monomeric form. Upon exposure to heat and

unactivated form. Binds to inverted 5'-NGAAN-3' pentamer DNA sequences. Binds to chromatin at heat shock gene promoters. Activates transcription of transcription factor FOXR1 which in turn activates transcription of the heat shock chaperones HSPA1A and HSPA6 and the antioxidant NADPH-dependent reductase DHRS2 (By similarity). Binds the promoter region upstream of exon 1 of Mpv17I to activate expression of the M-LPS isoform which is involved in metabolism of reactive oxygen species (PubMed:20231359). Also serves several other functions independently of its transcriptional activity. Involved in the repression of Ras-induced transcriptional activation of the c-fos gene in heat-stressed cells. Positively regulates pre-mRNA 3'-end processing and polyadenylation of HSP70 mRNA upon heat-stressed cells in a symplekin (SYMPK)-dependent manner. Plays a role in nuclear export of stress-induced HSP70 mRNA. Plays a role in the regulation of mitotic progression. Also plays a role as a negative regulator of non-homologous end joining (NHEJ) repair activity in a DNA damage-dependent manner. Involved in stress-induced cancer cell proliferation in a IER5-dependent manner.

Molecular Weight: 57.2 kDa

UniProt: P38532

Application Details

Application Notes: We expect the protein to work for functional studies. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months