

Datasheet for ABIN7563221 Id2 Protein (AA 1-134) (His tag)



Overview

Quantity:	1 mg
Target:	ld2
Protein Characteristics:	AA 1-134
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Id2 protein is labelled with His tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB)
Product Details	
Purpose:	Custom-made recombinat Id2 Protein expressed in mammalien cells.
Sequence:	MKAFSPVRSV RKNSLSDHSL GISRSKTPVD DPMSLLYNMN DCYSKLKELV PSIPQNKKVT
	KMEILQHVID YILDLQIALD SHPTIVSLHH QRPGQNQASR TPLTTLNTDI SILSLQASEF
	PSELMSNDSK VLCG Sequence without tag. The proposed Purification-Tag is based on

Characteristics:

Key Benefits:

• Made to order protein - from design to production - by highly experienced protein experts.

experiences with the expression system, a different complexity of the protein could make

- · Protein expressed in mammalien cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.

another tag necessary. In case you have a special request, please contact us.

• State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

Grade:

custom-made

Target Details

Target: Id2

Alternative Name:

Id2 (Id2 Products)

Background:

DNA-binding protein inhibitor ID-2 (Inhibitor of DNA binding 2) (Inhibitor of differentiation 2),FUNCTION: Transcriptional regulator (lacking a basic DNA binding domain) which negatively regulates the basic helix-loop-helix (bHLH) transcription factors by forming heterodimers and inhibiting their DNA binding and transcriptional activity. Implicated in regulating a variety of cellular processes, including cellular growth, senescence, differentiation, apoptosis, angiogenesis, and neoplastic transformation. Inhibits skeletal muscle and cardiac myocyte differentiation. Regulates the circadian clock by repressing the transcriptional activator activity of the CLOCK-BMAL1 heterodimer. Restricts the CLOCK and BMAL1 localization to the cytoplasm. Plays a role in both the input and output pathways of the circadian clock: in the input component, is involved in modulating the magnitude of photic entrainment and in the output component, contributes to the regulation of a variety of liver clock-controlled genes involved in lipid metabolism. {ECO:0000269|PubMed:19740747,

ECO:0000269|PubMed:20861012}.

Molecular Weight:

15.0 kDa

UniProt:

P41136

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	12 months