

Datasheet for ABIN7563515

TRAF6 Protein (AA 1-530) (His tag)



\sim				
O_1	/ el	rVI	161	Λ

Quantity:	1 mg
Target:	TRAF6
Protein Characteristics:	AA 1-530
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TRAF6 protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

Product Details

Purpose:	Custom-made recombinat Traf6 Protein expressed in mammalien cells.
Sequence:	MSLLNCENSC GSSQSSSDCC AAMAASCSAA VKDDSVSGSA STGNLSSSFM EEIQGYDVEF
	DPPLESKYEC PICLMALREA VQTPCGHRFC KACIIKSIRD AGHKCPVDNE ILLENQLFPD
	NFAKREILSL TVKCPNKGCL QKMELRHLED HQVHCEFALV NCPQCQRPFQ KCQVNTHIIE
	DCPRRQVSCV NCAVSMAYEE KEIHDQSCPL ANIICEYCGT ILIREQMPNH YDLDCPTAPI
	PCTFSVFGCH EKMQRNHLAR HLQENTQLHM RLLAQAVHNV NLALRPCDAA SPSRGCRPED
	PNYEETIKQL ESRLVRQDHQ IRELTAKMET QSMYVGELKR TIRTLEDKVA EMEAQQCNGI
	YIWKIGNFGM HLKSQEEERP VVIHSPGFYT GRPGYKLCMR LHLQLPTAQR CANYISLFVH
	TMQGEYDSHL PWPFQGTIRL TILDQSEALI RQNHEEVMDA KPELLAFQRP TIPRNPKGFG
	YVTFMHLEAL RQGTFIKDDT LLVRCEVSTR FDMGGLRKEG FQPRSTDAGV Sequence without
	tag. The proposed Purification-Tag is based on experiences with the expression system, a
	different complexity of the protein could make another tag necessary. In case you have a

special request, please contact us. Characteristics: Key Benefits: Made to order protein - from design to production - by highly experienced protein experts. Protein expressed in mammalien cells and purified in one-step affinity chromatography · The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins. • State-of-the-art algorithm used for plasmid design (Gene synthesis). This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein. If you are not interested in a full length protein, please contact us for individual protein fragments. The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified. > 90 % as determined by Bis-Tris Page, Western Blot Purity: Grade: custom-made **Target Details** TRAF6 Target: Alternative Name: Traf6 (TRAF6 Products) Background: TNF receptor-associated factor 6 (EC 2.3.2.27) (E3 ubiquitin-protein ligase TRAF6) (RING-type E3 ubiquitin transferase TRAF6), FUNCTION: E3 ubiquitin ligase that, together with UBE2N and UBE2V1, mediates the synthesis of 'Lys-63'-linked-polyubiquitin chains conjugated to proteins, such as ECSIT, IKBKG, IRAK1, AKT1 and AKT2 (PubMed:15322147, PubMed:17633018). Also mediates ubiquitination of free/unanchored polyubiquitin chain that leads to MAP3K7 activation (By similarity). Leads to the activation of NF-kappa-B and JUN. Seems to also play a role in dendritic cells (DCs) maturation and/or activation (PubMed:14499111). Represses c-Myb-mediated transactivation, in B-lymphocytes (By similarity). Adapter protein that seems to

play a role in signal transduction initiated via TNF receptor, IL-1 receptor and IL-17 receptor

activation of adapter protein complex 1 (AP-1) and NF-kappa-B, in response to RANK-L

(PubMed:10421844, PubMed:10215628). Regulates osteoclast differentiation by mediating the

stimulation (PubMed:10421844, PubMed:17092936). Together with MAP3K8, mediates CD40

rarget Details		
	signals that activate ERK in B-cells and macrophages, and thus may play a role in the regulation	
	of immunoglobulin production (PubMed:12881420). Participates also in the TCR signaling by	
	ubiquitinating LAT (By similarity). {ECO:0000250 UniProtKB:Q9Y4K3,	
	ECO:0000269 PubMed:10215628, ECO:0000269 PubMed:10421844,	
	ECO:0000269 PubMed:12881420, ECO:0000269 PubMed:14499111,	
	ECO:0000269 PubMed:15322147, ECO:0000269 PubMed:17092936,	
	ECO:0000269 PubMed:17633018}.	
Molecular Weight:	60.1 kDa	
UniProt:	P70196	
Pathways:	NF-kappaB Signaling, TCR Signaling, TLR Signaling, Fc-epsilon Receptor Signaling Pathway,	
	Neurotrophin Signaling Pathway, Activation of Innate immune Response, Regulation of	
	Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Production of	
	Molecular Mediator of Immune Response, Tube Formation, Hepatitis C, Toll-Like Receptors	
	Cascades, Ubiquitin Proteasome Pathway	
Application Details		
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies	
	as well. As the protein has not been tested for functional studies yet we cannot offer a	
	guarantee though.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	The buffer composition is at the discretion of the manufacturer.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	-80 °C	
Storage Comment:	Store at -80°C.	
Expiry Date:	12 months	