

Datasheet for ABIN7563584

## PKC zeta Protein (AA 1-592) (His tag)



[Go to Product page](#)

### Overview

Quantity:	1 mg
Target:	PKC zeta (PRKCZ)
Protein Characteristics:	AA 1-592
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This PKC zeta protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS)

### Product Details

Purpose:	Custom-made recombinat Prkcz Protein expressed in mammalian cells.
Sequence:	<p>MPSRTDPKMD RSGGRVRLKA HYGGDILITS VDAMTTFKDL CEEVRDMCGL HQQHPLTLKW</p> <p>VDSEGDPCVT SSQMELEEF RLVCQGRDEV LIHVFPSP EQPGMPCPG DSIYRRGAR</p> <p>RWRKLYRANG HLFQAKRFNR GAYCGQCSER IWGLSRQGYR CINCKLLVHK RCHVLVPLTC</p> <p>RRHMDSVMPS QEPPVDDKND GVDLPSEETD GIAYISSRK HDNIKDDSED LKPVIDGVDG</p> <p>IKISQGLGLQ DFDLIRVIGR GSYAKVLLVR LKKNDQIYAM KVVKKELVHD DEDIDWVQTE</p> <p>KHVFEQASSN PFLVGLHSCF QTTSRLFLVI EYVNGGDLMF HMQRQRKLPE EHARFYAAEI</p> <p>CIALNFLHER GIIYRDLKLD NVLLDADGHI KLTGYGMCKE GLGPGDTTST FCGTPNYIAP</p> <p>EILRGEEYGF SVDWWALGVL MFEMMAGRSP FDIITDNPDM NTEDYLFQVI LEKPIRIPRF</p> <p>LSVKASHVLK GFLNKDPKER LGCRPQTGFS DIKSHAFFRS IDWDLLEKKQ TLPPFQPQIT</p> <p>DDYGLDNFDT QFTSEPVQLT PDDEDVIKRI DQSEFEGFEY INPLLLSAEE SV <b>Sequence without tag.</b></p> <p><b>The proposed Purification-Tag is based on experiences with the expression system, a</b></p>

**different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.**

Characteristics:	<p>Key Benefits:</p> <ul style="list-style-type: none"><li>• Made to order protein - from design to production - by highly experienced protein experts.</li><li>• Protein expressed in mammalian cells and purified in one-step affinity chromatography</li><li>• The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.</li><li>• State-of-the-art algorithm used for plasmid design (Gene synthesis).</li></ul> <p>This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.</p> <p>If you are not interested in a full length protein, please contact us for individual protein fragments.</p> <p>The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.</p>
Purity:	> 90 % as determined by Bis-Tris Page, Western Blot
Grade:	custom-made

Target Details

Target:	PKC zeta (PRKCZ)
Alternative Name:	Prkcz ( <a href="#">PRKCZ Products</a> )
Background:	<p>Protein kinase C zeta type (EC 2.7.11.13) (nPKC-zeta),FUNCTION: Calcium- and diacylglycerol-independent serine/threonine-protein kinase that functions in phosphatidylinositol 3-kinase (PI3K) pathway and mitogen-activated protein (MAP) kinase cascade, and is involved in NF-kappa-B activation, mitogenic signaling, cell proliferation, cell polarity, inflammatory response and maintenance of long-term potentiation (LTP). Upon lipopolysaccharide (LPS) treatment in macrophages, or following mitogenic stimuli, functions downstream of PI3K to activate MAP2K1/MEK1-MAPK1/ERK2 signaling cascade independently of RAF1 activation. Required for insulin-dependent activation of AKT3, but may function as an adapter rather than a direct activator. Upon insulin treatment may act as a downstream effector of PI3K and contribute to the activation of translocation of the glucose transporter SLC2A4/GLUT4 and subsequent glucose transport in adipocytes. In EGF-induced cells, binds and activates MAP2K5/MEK5-</p>

## Target Details

MAPK7/ERK5 independently of its kinase activity and can activate JUN promoter through MEF2C. Through binding with SQSTM1/p62, functions in interleukin-1 signaling and activation of NF-kappa-B with the specific adapters RIPK1 and TRAF6. Participates in TNF-dependent transactivation of NF-kappa-B by phosphorylating and activating IKBKB kinase, which in turn leads to the degradation of NF-kappa-B inhibitors. In migrating astrocytes, forms a cytoplasmic complex with PARD6A and is recruited by CDC42 to function in the establishment of cell polarity along with the microtubule motor and dynein. In association with FEZ1, stimulates neuronal differentiation in PC12 cells. In the inflammatory response, is required for the T-helper 2 (Th2) differentiation process, including interleukin production, efficient activation of JAK1 and the subsequent phosphorylation and nuclear translocation of STAT6. May be involved in development of allergic airway inflammation (asthma), a process dependent on Th2 immune response. In the NF-kappa-B-mediated inflammatory response, can relieve SETD6-dependent repression of NF-kappa-B target genes by phosphorylating the RELA subunit at 'Ser-311'. Phosphorylates VAMP2 in vitro (By similarity). {ECO:0000250|UniProtKB:Q05513, ECO:0000269|PubMed:15987782, ECO:0000269|PubMed:21131967}., FUNCTION: [Isoform 2]: Involved in late synaptic long term potentiation phase in CA1 hippocampal cells and long term memory maintenance. {ECO:0000269|PubMed:27187150}.

Molecular Weight:	67.7 kDa
UniProt:	<a href="#">Q02956</a>
Pathways:	<a href="#">NF-kappaB Signaling</a> , <a href="#">RTK Signaling</a> , <a href="#">Myometrial Relaxation and Contraction</a> , <a href="#">Regulation of Leukocyte Mediated Immunity</a> , <a href="#">Positive Regulation of Immune Effector Process</a> , <a href="#">Synaptic Membrane</a> , <a href="#">Production of Molecular Mediator of Immune Response</a> , <a href="#">CXCR4-mediated Signaling Events</a> , <a href="#">Thromboxane A2 Receptor Signaling</a>

## Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.

Handling

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: 12 months