

Datasheet for ABIN7563617

NLR Family, Pyrin Domain Containing 1B (NLRP1B) (AA 1-1196) protein (His tag)



[Go to Product page](#)

Overview

Quantity:	1 mg
Target:	NLR Family, Pyrin Domain Containing 1B (NLRP1B)
Protein Characteristics:	AA 1-1196
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	His tag
Application:	SDS-PAGE (SDS), Western Blotting (WB)

Product Details

Purpose:	Custom-made recombinat Nlrp1b Protein expressed in mammalian cells.
Sequence:	MEESPPKQKS NTKVTQHEGQ QDLNTRHMN VELKHPKLE RHLKLGMPV VYMKQGEEIL YPAQSLREEN LIQNFTSLLL LQKLCPKDPE NMVRKSWASC VPEEGGHMIN IQDLFGPNIG TQKEPQLVII EGAAGIGKST LARLVKRAWK EGQLYRDHFQ HVFFFSCREL AQCKKLSLAE LIAQGQEVPT APINQILSHP EKLLFILDGI DEPAWVLADQ NPELCLHWSQ RQPVHTLLGS LLGKSIFPEA FFLLTTRTTA LQKFIPSLPM PCQVEVLGFS GIERENYFYK YFANQRHAIT AFMMVESNPV LLTLCEVPWV CWLVCTCLKK QMEQGRVLSL KSQTTTALCL KYLSLTIPDK HRRTQVKALC SLAAEGIWKR RTLFSESDLC KQGLDEDAVA TFLKTGVLQK QASSLSYSFA HLCLQEFAA ISCILEDSEE RHGNMEMDRI VETLVERYGR QNLFEAPT VR FLFGLLGKEG VKGMEKLFSC SLHGKTNLKL LWHILVKSQP HQPPCLGLLH CLYENQDMEL LTHVMHDLQG TIVPGPN DIA HTVLQTNVKH LVVQTDME LM VATFCIQFYC HVRTLQLNME KQQGYALISP RMVLYRWTP I TNASWEILFY NLKFTRNLEG LDLSGNSLRY SVVQSLCNTL RYPGCQLKTL

Product Details

WLVKCGLTSR HCSSLASVLS AHSSLTELYL QLNDLGDDGV RMLCEGLRNP VCNLSILWLD
LYLSAQVIT ELRTLEEKNP KLYIRSIWMP HMMVPTENMD EEAILTTFKQ QRQEPGDKPM
EILGTEEDFW GPTGPVATEL VDRVRNLYRV QLPMAGSYHC PSTGLHFVVT RAVTIEIEFC
AWSQFLDKTP LQQSHMVVGP LFDIKAEQGA VTAVYLPHFV SLKDTEASTF DFKVAHFQEH
GMVLETPDRV KPGYTVLKNP SFSPMGVVL R IIPAARHFIP ITSITLIYYR LNLEEVTLHL
YLVPNDCITQ KAIDDEEMKF QFVRINKPPP VDNLFIGSRY IVSGSENLEI TPKELELCYR
SSKEFQLFSE IYVGNMGSEI KLQIKNKKHM KLIWEALLKP EFKFDHLCDDQ EFCTILKNIM
ISPAGDLRPA LPRIAQALKD APSLLHFMDQ HREQLVARVT SVDPLLDKLH GLVLNEESYE
AVRAENTNQD KMRKLFNLSR SWSRACKDLF YQALKETHPH LVMDLFEKSG GVSLGS **Sequence
without tag. The proposed Purification-Tag is based on experiences with the expression
system, a different complexity of the protein could make another tag necessary. In case you
have a special request, please contact us.**

Characteristics:

Key Benefits:

- Made to order protein - from design to production - by highly experienced protein experts.
- Protein expressed in mammalian cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity: > 90 % as determined by Bis-Tris Page, Western Blot

Grade: custom-made

Target Details

Target: NLR Family, Pyrin Domain Containing 1B (NLRP1B)

Alternative Name: Nlrp1b ([NLRP1B Products](#))

Background: NACHT, LRR and PYD domains-containing protein 1b allele 5 (EC 3.4.-.-) [Cleaved into: NACHT,

LRR and PYD domains-containing protein 1b, C-terminus (Nlrp1b1-CT), NACHT, LRR and PYD domains-containing protein 1b, N-terminus (Nlrp1b1-NT)],FUNCTION: Acts as the sensor component of the Nlrp1b inflammasome, which mediates inflammasome activation in response to various pathogen-associated signals, leading to subsequent pyroptosis (By similarity). Inflammasomes are supramolecular complexes that assemble in the cytosol in response to pathogens and other damage-associated signals and play critical roles in innate immunity and inflammation (By similarity). Acts as a recognition receptor (PRR): recognizes specific pathogens and other damage-associated signals, such as B.anthraxis lethal toxin (LT) or Val-boroPro inhibitor, and mediates the formation of the inflammasome polymeric complex (PubMed:16429160, PubMed:31383852). In response to pathogen-associated signals, the N-terminal part of Nlrp1b is degraded by the proteasome, releasing the cleaved C-terminal part of the protein (NACHT, LRR and PYD domains-containing protein 1b, C-terminus), which polymerizes to initiate the formation of the inflammasome complex: the inflammasome directly recruits pro-caspase-1 (proCASP1) independently of PYCARD/ASC and promotes caspase-1 (CASP1) activation, which subsequently cleaves and activates inflammatory cytokines IL1B and IL18 and gasdermin-D (GSDMD), leading to pyroptosis (By similarity). In the absence of GSDMD expression, the Nlrp1b inflammasome is able to recruit and activate CASP8, leading to activation of gasdermin-E (GSDME) (By similarity). Activation of Nlrp1b inflammasome is also required for HMGB1 secretion, the active cytokines and HMGB1 stimulate inflammatory responses (By similarity). Primary mediator of macrophage susceptibility to B.anthraxis LT: in response to B.anthraxis infection, macrophages and dendritic cells release IL1B and undergo pyroptosis (PubMed:16429160, PubMed:24935976). This early inflammatory response to the toxin increases resistance to infection by B.anthraxis spores (PubMed:16429160, PubMed:24935976). {ECO:0000250|UniProtKB:Q2LKW6, ECO:0000269|PubMed:16429160, ECO:0000269|PubMed:24935976, ECO:0000269|PubMed:31383852}., FUNCTION: [NACHT, LRR and PYD domains-containing protein 1b allele 5]: Constitutes the precursor of the Nlrp1b inflammasome, which mediates autoproteolytic processing within the FIIND domain to generate the N-terminal and C-terminal parts, which are associated non-covalently in absence of pathogens and other damage-associated signals. {ECO:0000250|UniProtKB:Q2LKW6}., FUNCTION: [NACHT, LRR and PYD domains-containing protein 1b, N-terminus]: Regulatory part that prevents formation of the Nlrp1b inflammasome: in absence of pathogens and other damage-associated signals, interacts with the C-terminal part of Nlrp1b (NACHT, LRR and PYD domains-containing protein 1b, C-terminus), preventing activation of the Nlrp1b inflammasome. In response to pathogen-associated signals, this part is ubiquitinated by the N-end rule pathway and degraded by the proteasome, releasing the cleaved C-terminal part of the protein, which polymerizes and forms the Nlrp1b inflammasome. {ECO:0000250|UniProtKB:Q2LKW6}.,

Target Details

FUNCTION: [NACHT, LRR and PYD domains-containing protein 1b, C-terminus]: Constitutes the active part of the Nlrp1b inflammasome. In absence of pathogens and other damage-associated signals, interacts with the N-terminal part of Nlrp1b (NACHT, LRR and PYD domains-containing protein 1b, N-terminus), preventing activation of the Nlrp1b inflammasome. In response to pathogen-associated signals, the N-terminal part of Nlrp1b is degraded by the proteasome, releasing this form, which polymerizes to form the Nlrp1b inflammasome complex: the Nlrp1b inflammasome complex then directly recruits pro-caspase-1 (proCASP1) and promotes caspase-1 (CASP1) activation, leading to gasdermin-D (GSDMD) cleavage and subsequent pyroptosis. {ECO:0000250|UniProtKB:Q2LKW6}.

Molecular Weight: 136.5 kDa

UniProt: [Q0GKD5](#)

Application Details

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: The buffer composition is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: 12 months