

Datasheet for ABIN7563723

BRISC and BRCA1 A Complex Member 1 (BABAM1) (AA 1-333) protein (His tag)



Go to Product page

	er		

Characteristics:

1 mg
BRISC and BRCA1 A Complex Member 1 (BABAM1)
AA 1-333
Mouse
HEK-293 Cells
Recombinant
His tag
Western Blotting (WB), SDS-PAGE (SDS)
Custom-made recombinat Babam1 Protein expressed in mammalien cells.
MEVAEANSPT EEEEEEEEG EETISEPRPH TRSNPEGAED RALGAQASVG SRSEGEGEAA
TADGGAASVP GAGPKPWQVP ASASEVQIRT PRVNCPEKVI ICLDLSEEMS VPKLESFNGS
RTNALNVSQK MVEMFVRTKH KIDKSHEFAL VVVNDDSAWL SGLTSDPREL CSCLYDLETA
SCSTFNLEGL FSLIQQKTEL PVTENVQTIP PPYVVRTILV YSRPPCQPQF SLTEPMKKMF
QCPYFFFDIV YIHNGTEEKE EDMSWKDMFA FMGSLDTKGA SYKYEVALAG PALELHNCMA
KLLAHPLQRP CQTHASYSLL EEDEEAGEEE ATV Sequence without tag. The proposed
Purification-Tag is based on experiences with the expression system, a different complexity
• • • • • • • • • • • • • • • • • • •

contact us.

Key Benefits:

- · Made to order protein from design to production by highly experienced protein experts.
- · Protein expressed in mammalien cells and purified in one-step affinity chromatography
- The optimized expression system ensures reliability for intracellular, secreted and transmembrane proteins.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab try to ensure that you receive soluble protein.

If you are not interested in a full length protein, please contact us for individual protein fragments.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

Purity:

> 90 % as determined by Bis-Tris Page, Western Blot

Grade:

custom-made

Target Details

Target:

BRISC and BRCA1 A Complex Member 1 (BABAM1)

Alternative Name:

Babam1 (BABAM1 Products)

Background:

BRISC and BRCA1-A complex member 1 (Mediator of RAP80 interactions and targeting subunit of 40 kDa) (New component of the BRCA1-A complex), FUNCTION: Component of the BRCA1-A complex, a complex that specifically recognizes 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA lesions sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at double-strand breaks (DSBs). The BRCA1-A complex also possesses deubiquitinase activity that specifically removes 'Lys-63'-linked ubiquitin on histones H2A and H2AX. In the BRCA1-A complex, it is required for the complex integrity and its localization at DSBs.

Component of the BRISC complex, a multiprotein complex that specifically cleaves 'Lys-63'-linked ubiquitin in various substrates. In these 2 complexes, it is probably required to maintain the stability of BABAM2 and help the 'Lys-63'-linked deubiquitinase activity mediated by BRCC3/BRCC36 component. The BRISC complex is required for normal mitotic spindle assembly and microtubule attachment to kinetochores via its role in deubiquitinating NUMA1. Plays a role in interferon signaling via its role in the deubiquitination of the interferon receptor IFNAR1, deubiquitination increases IFNAR1 activity by enhancing its stability and cell surface

Target Details

Expiry Date:

12 months

Target Details	
	expression. Down-regulates the response to bacterial lipopolysaccharide (LPS) via its role in
	IFNAR1 deubiquitination. {ECO:0000250 UniProtKB:Q9NWV8}.
Molecular Weight:	36.8 kDa
UniProt:	Q3UI43
Pathways:	Positive Regulation of Response to DNA Damage Stimulus
Application Details	
Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies
	as well. As the protein has not been tested for functional studies yet we cannot offer a
	guarantee though.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.